

Exercises discussed on November 6, 2012

Recall the definition of *Stirling numbers of the second kind* $S_2(n, k)$ as the number of ways to partition an n -element set into a disjoint union of k nonempty subsets. They satisfy the recurrence relation

$$S_2(n, k) = S_2(n-1, k-1) + kS_2(n-1, k), \quad n, k \geq 1,$$

with initial values $S_2(0, 0) = 1$ and $S_2(n, 0) = 0$ for $n \geq 1$, and $S_2(n, k) = 0$ for $k > n \geq 1$.

19. Show that for $k \in \mathbb{N}$

$$\sum_{n=0}^{\infty} S_2(n, k)x^n = \frac{x^k}{(1-x)(1-2x)\cdots(1-kx)}.$$

20. Show that for $k \in \mathbb{N}$

$$\sum_{n=0}^{\infty} S_2(n, k) \frac{x^n}{n!} = \frac{1}{k!} (e^x - 1)^k,$$

and that

$$\sum_{n,k=0}^{\infty} S_2(n, k) \frac{x^n}{n!} y^k = \exp(y(e^x - 1)).$$

21. Let the signless Stirling numbers of the first kind $C(n, k)$ denote the number of permutations of $\{1, 2, \dots, n\}$ with exactly k cycles. Derive a recurrence relation for $C(n, k)$. Starting from this recurrence, derive a recurrence relation for the Stirling numbers of the first kind $S_1(n, k) := (-1)^{n-k} C(n, k)$.

22. Let x be an indeterminate and $n \in \mathbb{N}$. Show that

(a) $x^n = \sum_{k=0}^n S_2(n, k)x^k$

(b) $x^n = \sum_{k=0}^n S_1(n, k)x^k$