

# Introduction to Unification Theory

## Higher-Order Unification

Temur Kutsia

RISC, Johannes Kepler University of Linz, Austria  
`kutsia@risc.uni-linz.ac.at`



# Overview

Introduction

Preliminaries

Higher-Order Unification Procedure

# Outline

Introduction

Preliminaries

Higher-Order Unification Procedure

# Introduction

- ▶ In first order unification, we were not allowed to replace a variable with a function.

# Introduction

- ▶ In first order unification, we were not allowed to replace a variable with a function.
- ▶ However, it makes sense to ask to find, e.g., a function that when applied to an object gives again this object: Find an  $F$  such that  $F(a) = a$ .



# Introduction

- ▶ In first order unification, we were not allowed to replace a variable with a function.
- ▶ However, it makes sense to ask to find, e.g., a function that when applied to an object gives again this object: Find an  $F$  such that  $F(a) = a$ .
- ▶  $F$ : Higher-order variable, appears at functional position.



# Introduction

- ▶ In first order unification, we were not allowed to replace a variable with a function.
- ▶ However, it makes sense to ask to find, e.g., a function that when applied to an object gives again this object: Find an  $F$  such that  $F(a) = a$ .
- ▶  $F$ : Higher-order variable, appears at functional position.
- ▶ Can be solved, e.g., with the identity function or with the constant function  $a$ .



# Introduction

- ▶ In first order unification, we were not allowed to replace a variable with a function.
- ▶ However, it makes sense to ask to find, e.g., a function that when applied to an object gives again this object: Find an  $F$  such that  $F(a) = a$ .
- ▶  $F$ : Higher-order variable, appears at functional position.
- ▶ Can be solved, e.g., with the identity function or with the constant function  $a$ .
- ▶ Higher-order equations.





# Introduction

- ▶ In first order unification, we were not allowed to replace a variable with a function.
- ▶ However, it makes sense to ask to find, e.g., a function that when applied to an object gives again this object: Find an  $F$  such that  $F(a) = a$ .
- ▶  $F$ : Higher-order variable, appears at functional position.
- ▶ Can be solved, e.g., with the identity function or with the constant function  $a$ .
- ▶ Higher-order equations.
- ▶ Solving method: Higher-order unification.



# Introduction

- ▶ Higher-order unification is fundamental in automating higher-order reasoning.
- ▶ Used in logical frameworks, logic programming, program synthesis, program transformation, type inferencing, computational linguistics, etc.
- ▶ Much more complicated than first-order unification (undecidable, of type zero, nonterminating, ...).
- ▶ In this lecture: Introduction to higher-order unification.



# Outline

Introduction

**Preliminaries**

Higher-Order Unification Procedure

# Simply Typed $\lambda$ -Calculus

- ▶ Simply type  $\lambda$ -calculus is our term language.
- ▶ In this section: Definitions and elementary properties.
  - ▶ Types
  - ▶ Terms
  - ▶ Substitutions
  - ▶ Reduction
  - ▶ Unification



# Types

## Types

Consider a finite set whose elements are called *atomic types* (or *base types*). Then:

- ▶ Atomic types are types,
- ▶ If  $T$  and  $U$  are types then  $T \rightarrow U$  is a type.

The expression  $T_1 \rightarrow T_2 \rightarrow \cdots \rightarrow T_n \rightarrow U$  is a notation for the type  $T_1 \rightarrow (T_2 \rightarrow \cdots \rightarrow (T_n \rightarrow U) \dots)$ .



# Types

## Order of a Type

- ▶  $o(T) = 1$  if  $T$  is atomic.
- ▶  $o(T \rightarrow U) = \max\{1 + o(T), o(U)\}$ .

## Example

Let  $T_1, T_2, T_3$  be atomic types, then

- ▶  $o(T_1 \rightarrow T_2 \rightarrow T_3) = 2$ .
- ▶  $o((T_1 \rightarrow T_2) \rightarrow T_3) = 3$ .



# Terms

Assumptions:

- ▶ Consider finite set of constants.
- ▶ To each constant a type is assigned.
- ▶ For each atomic type there is at least one constant.
- ▶ For each type there is an infinite set of variables.
- ▶ Two different types have disjoint sets of variables.

## $\lambda$ -Terms

- ▶ Constants are terms.
- ▶ Variables are terms.
- ▶ If  $t$  and  $s$  are terms then  $(t s)$  is a term.
- ▶ If  $x$  is a variable and  $t$  is a term then  $\lambda x. t$  is a term.

The expression  $(t s_1 \dots s_n)$  is a notation for the term  
 $(\dots (t s_1) \dots s_n)$



# Terms

- ▶  $\lambda x. t$  is a function where  $\lambda x$  is the  $\lambda$ -abstraction and  $t$  is the body. Intuitively, it is a function  $x \mapsto t$ .
- ▶ In  $\lambda x. t$ ,  $\lambda x$  is a binder for  $x$  in  $t$ . Occurrences of  $x$  in  $t$  are *bound*.
- ▶  $(t s)$  is an application where function  $t$  is applied to the argument  $s$ .





# Terms

## Type of a Term

A term  $t$  is said to have the type  $T$  if either

- ▶  $t$  is a constant of type  $T$ ,
  - ▶  $t$  is a variable of type  $T$ ,
  - ▶  $t = (r s)$ ,  $r$  has type  $U \rightarrow T$  and  $s$  has type  $U$  for some  $U$ ,
  - ▶  $t = \lambda x. s$ , the variable  $x$  has type  $U$ , the term  $s$  has type  $V$  and  $T = U \rightarrow V$ .
- 
- ▶ A term  $t$  is said to be *well-typed* if there exists a type  $T$  such that  $t$  has type  $T$ .
  - ▶ In this case  $T$  is unique and it is called *the type of  $t$* .
  - ▶ We consider only well-typed terms.



# Order

## Order of a Symbol, Language

- ▶ The order of a function symbol or a variable is the order of its type.
- ▶ A language of order  $n$  is one which allows function symbols of order at most  $n + 1$  and variables of order at most  $n$ .

Formalization of the conventions:

- ▶ First order term denotes an individual.
- ▶ Second order term denotes a function on individuals.
- ▶ etc.



# Free Variables

- ▶  $vars(t)$ : The set of variables occurring in the term  $t$ .
- ▶ An occurrence of a variable in a term is *free* if it is not bound.
- ▶ The set of variables that occur freely in  $t$ , denoted  $fvars(t)$ :
  - ▶  $fvars(c) = \emptyset$ , where  $c$  is a constant.
  - ▶  $fvars(x) = \{x\}$ .
  - ▶  $fvars((s r)) = fvars(s) \cup fvars(r)$ .
  - ▶  $fvars(\lambda x. s) = fvars(s) \setminus \{x\}$ .
- ▶ Closed term: A term without free variables.



# Free Variables

## Example

- ▶  $fvars(\lambda x. x) = \emptyset$ .  
(Closed term)
- ▶  $fvars(\lambda x. y) = \{y\}$ .
- ▶  $fvars(((\lambda x. x) x)) = \{x\}$ .  
( $x$  has a bound occurrence as well)



# Substitution

- ▶ We reuse the definition of substitution as finite mapping from the previous lectures, but in addition require that it preserves types.
- ▶ Hence, if  $x \mapsto t$  is a binding of a substitution,  $x$  and  $t$  have the same type.
- ▶ The definitions of composition, more general substitution, etc. will also be reused.



# Replacement in a Term

## Replacement in a Term

Let  $\sigma = \{x_1 \mapsto t_1, \dots, x_n \mapsto t_n\}$  be a substitution and  $t$  be a term, then the term  $t\langle\sigma\rangle$  is defined as follows:

- ▶  $c\langle\sigma\rangle = c$ .
- ▶  $x_i\langle\sigma\rangle = t_i$ .
- ▶  $x\langle\sigma\rangle = x$ , if  $x \notin \{x_1, \dots, x_n\}$ .
- ▶  $(sr)\langle\sigma\rangle = (s\langle\sigma\rangle r\langle\sigma\rangle)$ .
- ▶  $(\lambda x. s)\langle\sigma\rangle = (\lambda x. s\langle\sigma\rangle)$ .

## Example

- ▶  $(\lambda x. x)\langle\{x \mapsto y\}\rangle = \lambda x. y$ .
- ▶  $(\lambda y. x)\langle\{x \mapsto y\}\rangle = \lambda y. y$  (variable capture).



# $\alpha$ -Equivalence

## $\alpha$ -Equivalence

- ▶  $c \equiv_{\alpha} c.$
- ▶  $x \equiv_{\alpha} x.$
- ▶  $(ts) \equiv_{\alpha} (t's')$  if  $t \equiv_{\alpha} t'$  and  $s \equiv_{\alpha} s'$ .
- ▶  $\lambda x. t \equiv_{\alpha} \lambda y. s$  if  $t\langle\{x \mapsto z\}\rangle \equiv_{\alpha} s\langle\{y \mapsto z\}\rangle$  for some variable  $z$  different from  $x$  and  $y$  and occurring neither in  $t$  nor in  $s$ .

## Example

- ▶  $\lambda x. x \equiv_{\alpha} \lambda y. y.$
- ▶  $\alpha$ -equivalence is an equivalence relation.
- ▶ Application and abstraction are compatible with  $\alpha$ -equivalence.



# Substitution in a Term

## Substitution in a Term

Let  $\sigma = \{x_1 \mapsto t_1, \dots, x_n \mapsto t_n\}$  be a substitution and  $t$  be a term, then the term  $t\sigma$  is defined as follows:

- ▶  $c\sigma = c$ .
- ▶  $x_i\sigma = t_i$ .
- ▶  $x\sigma = x$ , if  $x \notin \{x_1, \dots, x_n\}$ .
- ▶  $(sr)\sigma = (s\sigma r\sigma)$ .
- ▶  $(\lambda x. s)\sigma = (\lambda y. s\{x \mapsto y\}\sigma)$ , where  $y$  is a fresh variable of the same type as  $x$ .

Since the choice of fresh variable is arbitrary, the substitution operation is defined on  $\alpha$ -equivalence classes.





# Substitution in a Term

## Example

- ▶  $(\lambda x. x)\{x \mapsto y\} = \lambda z. z.$
- ▶  $(\lambda y. x)\{x \mapsto y\} = \lambda z. y$  (no variable capture).
- ▶  $(x \lambda x. (x y))\{x \mapsto \lambda z. z\} = (\lambda z. z \lambda u. (u y)).$

# Reduction

- ▶ Intuition: Function evaluation.
- ▶ For instance, evaluating function  $f : x \mapsto x + 1$  at 2:  
 $f(2) = 2 + 1$ .
- ▶ As  $\lambda$ -terms:  $((\lambda x. x + 1) 2) \triangleright x + 1\{x \mapsto 2\} = 2 + 1$ .  
( $\beta$ -reduction)



# Reduction

Formally:

## $\beta\eta$ -Reduction

- ▶  $\beta$ -reduction:  $((\lambda x.s) t) \triangleright s\{x \mapsto t\}$ .
- ▶  $\eta$ -reduction:  $(\lambda x.(t x)) \triangleright t$ , if  $x \notin fvars(t)$ .

Propagates into contexts:

- ▶ If  $s \triangleright s'$  then  $(s t) \triangleright (s' t)$ .
- ▶ If  $t \triangleright t'$  then  $(s t) \triangleright (s t')$ .
- ▶ If  $t \triangleright t'$  then  $\lambda x. t \triangleright \lambda x. t'$ .



# Reduction

$\triangleright^*$  - reflexive-transitive closure of  $\triangleright$ .

Facts:

- ▶  $\beta\eta$ -Reduction preserves types.
- ▶ If  $s \triangleright^* t$  then  $s\sigma \triangleright^* t\sigma$ .
- ▶ Each term has a unique  $\beta\eta$ -normal form modulo  $\alpha$ -equivalence.



# Reduction

## Example

$$\begin{aligned}\lambda x.(f((\lambda y.(y x)) \lambda z.z)) &\triangleright_{\beta} \lambda x.(f((\lambda z.z) x)) \\ &\triangleright_{\beta} \lambda x.(f x) \\ &\triangleright_{\eta} f\end{aligned}$$

# Long Normal Form

## Long Normal Form

Assume

- ▶  $t = \lambda x_1 \dots \lambda x_m. (r s_1 \dots s_k)$  is in the  $\beta\eta$ -normal form,
- ▶  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  is a type of  $t$ ,
- ▶  $U$  is atomic and  $n \geq m$ .

Then the long normal form of  $t$  is the term

$$t' = \lambda x_1 \dots \lambda x_m. \lambda x_{m+1} \dots \lambda x_n. (r s'_1 \dots s'_k x'_{m+1} \dots x'_n)$$

where

- ▶  $s'_j$  is the long normal form of  $s_j$ .
- ▶  $x'_j$  is the long normal form of  $x_j$ .



# Long Normal Form

## Long Normal Form

Assume

- ▶  $t = \lambda x_1 \dots \lambda x_m. (r s_1 \dots s_k)$  is in the  $\beta\eta$ -normal form,
- ▶  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  is a type of  $t$ ,
- ▶  $U$  is atomic and  $n \geq m$ .

Then the long normal form of  $t$  is the term

$$t' = \lambda x_1 \dots \lambda x_m. \lambda x_{m+1} \dots \lambda x_n. (r s'_1 \dots s'_k x'_{m+1} \dots x'_n)$$

where

- ▶  $s'_j$  is the long normal form of  $s_j$ .
- ▶  $x'_j$  is the long normal form of  $x_j$ .

The long normal form of any term is that of its normal form.



# Long Normal Form

## Long Normal Form

Assume

- ▶  $t = \lambda x_1 \dots \lambda x_m. (r s_1 \dots s_k)$  is in the  $\beta\eta$ -normal form,
- ▶  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  is a type of  $t$ ,
- ▶  $U$  is atomic and  $n \geq m$ .

Then the long normal form of  $t$  is the term

$$t' = \lambda x_1 \dots \lambda x_m. \lambda x_{m+1} \dots \lambda x_n. (r s'_1 \dots s'_k x'_{m+1} \dots x'_n)$$

where

- ▶  $s'_j$  is the long normal form of  $s_j$ .
- ▶  $x'_j$  is the long normal form of  $x_j$ .

The long normal form of any term is that of its normal form.

Since  $t$  is in the normal form,  $r$  (called the *head* of  $t$ ) is either a constant or a variable.





# Long Normal Form

## Example

Let the type of  $f$  be  $T_1 \rightarrow T_2 \rightarrow U$ , with  $U$  atomic.

Let  $t$  be  $\lambda x.(f((\lambda y.(y\ x))\ \lambda z.z))$ .



# Long Normal Form

## Example

Let the type of  $f$  be  $T_1 \rightarrow T_2 \rightarrow U$ , with  $U$  atomic.

Let  $t$  be  $\lambda x.(f((\lambda y.(y\ x))\ \lambda z.z))$ .

- ▶ The long normal form of  $t$  is  $\lambda x.\lambda y.(f\ x\ y)$ .



# Long Normal Form

## Example

Let the type of  $f$  be  $T_1 \rightarrow T_2 \rightarrow U$ , with  $U$  atomic.

Let  $t$  be  $\lambda x.(f((\lambda y.(y\ x))\ \lambda z.z))$ .

- ▶ The long normal form of  $t$  is  $\lambda x.\lambda y.(f\ x\ y)$ .
- ▶  $\lambda x.\lambda y.(f\ x\ y)$  is a long normal form of  $\lambda x.(f\ x)$  as well, which is a  $\beta$ -normal form of  $t$ .



# Long Normal Form

## Example

Let the type of  $f$  be  $T_1 \rightarrow T_2 \rightarrow U$ , with  $U$  atomic.

Let  $t$  be  $\lambda x.(f((\lambda y.(y x)) \lambda z.z))$ .

- ▶ The long normal form of  $t$  is  $\lambda x.\lambda y.(f x y)$ .
- ▶  $\lambda x.\lambda y.(f x y)$  is a long normal form of  $\lambda x.(f x)$  as well, which is a  $\beta$ -normal form of  $t$ .
- ▶ In general, to compute long normal form, it is not necessary to perform  $\eta$ -reductions.



# Long Normal Form

- ▶ In the rest, “normal form” stands for “long normal form”.
- ▶ Notation: We write

$$\lambda x_1 \dots \lambda x_n. r(t_1, \dots, t_m)$$

for

$$\lambda x_1 \dots \lambda x_n. (r t_1 \dots t_m)$$

in normal form.  $r$  is either a constant or a variable.



# Complete Set of Unifiers

- ▶ A substitution  $\sigma$  is *more general on a set of variables  $\mathcal{X}$*  than  $\vartheta$ , written  $\sigma \leq^{\mathcal{X}} \vartheta$ , if there exists  $\psi$  such that  $x\sigma\psi \doteq x\vartheta$  for all  $x \in \mathcal{X}$  (modulo  $\beta\eta$ -reduction).



## Complete Sets of Unifiers

- ▶  $\mathcal{C}$  is a *complete set of unifiers* of a unification problem  $\Gamma$  iff
  1.  $\mathcal{C}$ 's elements are unifiers of  $\Gamma$ , and
  2. For each unifier of  $\Gamma$ ,  $\vartheta$ , there exists  $\sigma \in \mathcal{C}$  such that  $\sigma \leq^{\mathcal{X}} \vartheta$ , where  $\mathcal{X} = fvars(\Gamma)$ .
- ▶  $\mathcal{C}$  is a *minimal complete set of unifiers* of  $\Gamma$  if it is a complete set of unifiers of  $\Gamma$  and
  3. two distinct elements of  $\mathcal{C}$  are not comparable wrt  $\leq^{\mathcal{X}}$ .



# Outline

Introduction

Preliminaries

Higher-Order Unification Procedure



# Higher Order Unification

## Higher-Order Unification Problem, Unifier

- ▶ Higher-Order Unification problem: a finite set of equations

$$\Gamma = \{s_1 \doteq? t_1, \dots, s_n \doteq? t_n\},$$

where  $s_i, t_i$  are  $\lambda$ -terms.

- ▶ Unifier of  $\Gamma$ : a substitution  $\sigma$  such that  $s_i\sigma$  and  $t_i\sigma$  have the same normal form for each  $1 \leq i \leq n$ .

We will use capital letters to denote free variables in unification problems.





# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}.$

# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Unifier:  $\sigma_1 = \{F \mapsto \lambda x.f(x, b)\}$ .



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Unifier:  $\sigma_1 = \{F \mapsto \lambda x.f(x, b)\}$ .
- ▶ Justification:

$$F(f(a, b))\sigma_1 = ((\lambda x.f(x, b)) f(a, b)) \triangleright_{\beta} f(f(a, b), b).$$



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Unifier:  $\sigma_1 = \{F \mapsto \lambda x.f(x, b)\}$ .
- ▶ Justification:

$$F(f(a, b))\sigma_1 = ((\lambda x.f(x, b)) f(a, b)) \triangleright_{\beta} f(f(a, b), b).$$

$$f(F(a), b)\sigma_1 = f(((\lambda x.f(x, b)) a), b) \triangleright_{\beta} f(f(a, b), b).$$



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}.$

# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Another unifier:  $\sigma_2 = \{F \mapsto \lambda x.f(f(x, b), b)\}$ .



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Another unifier:  $\sigma_2 = \{F \mapsto \lambda x.f(f(x, b), b)\}$ .
- ▶ Justification:

$$F(f(a, b))\sigma_2 = ((\lambda x.f(f(x, b), b)) f(a, b)) \triangleright_{\beta} f(f(f(a, b), b), b).$$



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq^? f(F(a), b)\}$ .
- ▶ Another unifier:  $\sigma_2 = \{F \mapsto \lambda x.f(f(x, b), b)\}$ .
- ▶ Justification:

$$F(f(a, b))\sigma_2 = ((\lambda x.f(f(x, b), b)) f(a, b)) \triangleright_{\beta} f(f(f(a, b), b), b).$$

$$f(F(a), b)\sigma_2 = f(((\lambda x.f(f(x, b), b)) a), b) \triangleright_{\beta} f(f(f(a, b), b), b).$$





# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}.$

# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Infinitely many unifiers, of the shape

$$\{F \mapsto \lambda x. f(\dots f(x, b), \dots b)\}.$$



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Infinitely many unifiers, of the shape

$$\{F \mapsto \lambda x. f(\dots f(x, b), \dots b)\}.$$

- ▶ Incomparable wrt instantiation quasi-ordering.



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Infinitely many unifiers, of the shape

$$\{F \mapsto \lambda x. f(\dots f(x, b), \dots b)\}.$$

- ▶ Incomparable wrt instantiation quasi-ordering.
- ▶ Minimal complete set of unifiers.



# Higher Order Unification

## Example

- ▶  $\Gamma = \{F(f(a, b)) \doteq? f(F(a), b)\}$ .
- ▶ Infinitely many unifiers, of the shape

$$\{F \mapsto \lambda x. f(\dots f(x, b), \dots b)\}.$$

- ▶ Incomparable wrt instantiation quasi-ordering.
- ▶ Minimal complete set of unifiers.
- ▶ There are problems for which this set does not exist!



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .

# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions (together with the **instance terms**):

# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions (together with the **instance terms**):

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x) \} \quad H(\lambda x. G(x))$$





# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions (together with the **instance terms**):

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x) \quad H(\lambda x. G(x))$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y \quad Y$$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq^? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions (together with the **instance terms**):

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x)\} \quad H(\lambda x. G(x))$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y\} \quad Y$$

$$\sigma_1 = \{F \mapsto \lambda x. \lambda y. G_1(x, x(H_1^1(x, y))), G \mapsto \lambda x. Y\} \quad G_1(\lambda x. Y, Y)$$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq^? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions (together with the **instance terms**):

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x)\} \quad H(\lambda x. G(x))$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y\} \quad Y$$

$$\sigma_1 = \{F \mapsto \lambda x. \lambda y. G_1(x, x(H_1^1(x, y))), G \mapsto \lambda x. Y\} \quad G_1(\lambda x. Y, Y)$$

$$\sigma_2 = \{F \mapsto \lambda x. \lambda y. G_2(x, x(H_1^2(x, y)), x(H_2^2(x, y))), G \mapsto \lambda x. Y\} \\ G_2(\lambda x. Y, Y, Y)$$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions (together with the **instance terms**):

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x)\} \quad H(\lambda x. G(x))$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y\} \quad Y$$

$$\sigma_1 = \{F \mapsto \lambda x. \lambda y. G_1(x, x(H_1^1(x, y))), G \mapsto \lambda x. Y\} \quad G_1(\lambda x. Y, Y)$$

$$\sigma_2 = \{F \mapsto \lambda x. \lambda y. G_2(x, x(H_1^2(x, y)), x(H_2^2(x, y))), G \mapsto \lambda x. Y\} \\ G_2(\lambda x. Y, Y, Y)$$

...

$$\sigma_n = \{F \mapsto \lambda x. \lambda y. G_n(x, x(H_1^n(x, y)), \dots, x(H_n^n(x, y))), G \mapsto \lambda x. Y\} \\ G_n(\lambda x. Y, Y, \dots, Y) \quad (n \text{ } Y\text{'s})$$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions:

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x)\}$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y\}$$

$$\sigma_n = \{F \mapsto \lambda x. \lambda y. G_n(x, x(H_1^n(x, y)), \dots, x(H_n^n(x, y))), G \mapsto \lambda x. Y\}$$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq^? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions:

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x)\}$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y\}$$

$$\sigma_n = \{F \mapsto \lambda x. \lambda y. G_n(x, x(H_1^n(x, y)), \dots, x(H_n^n(x, y))), G \mapsto \lambda x. Y\}$$

- ▶ No minimal complete set of unifiers.

For all  $i, j > i$ :  $\sigma_i \not\leq^{\{F, G\}} \sigma_j$ ,  $\sigma \not\leq^{\{F, G\}} \sigma_i$ ,  $\sigma_i \not\leq^{\{F, G\}} \sigma$ , and  $\sigma_i = \{F, G\} \sigma_{i+1} \vartheta_i$  where

$$\vartheta_i = \{G_{i+1} \mapsto \lambda x. \lambda y_1. \dots \lambda y_{i+1}. G_i(x, y_1, \dots, y_i), \\ H_1^{i+1} \mapsto H_1^i, \dots, H_i^{i+1} \mapsto H_i^i\}$$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ Complete set of solutions:

$$\sigma = \{F \mapsto \lambda x. \lambda y. H(x)\}$$

$$\sigma_0 = \{F \mapsto \lambda x. x, G \mapsto \lambda x. Y\}$$

$$\sigma_n = \{F \mapsto \lambda x. \lambda y. G_n(x, x(H_1^n(x, y)), \dots, x(H_n^n(x, y))), G \mapsto \lambda x. Y\}$$

- ▶ No minimal complete set of unifiers.

For all  $i, j > i$ :  $\sigma_i \not\leq^{\{F, G\}} \sigma_j$ ,  $\sigma \not\leq^{\{F, G\}} \sigma_i$ ,  $\sigma_i \not\leq^{\{F, G\}} \sigma$ , and  $\sigma_i =^{\{F, G\}} \sigma_{i+1} \vartheta_i$  where

$$\vartheta_i = \{G_{i+1} \mapsto \lambda x. \lambda y_1. \dots \lambda y_{i+1}. G_i(x, y_1, \dots, y_i), \\ H_1^{i+1} \mapsto H_1^i, \dots, H_i^{i+1} \mapsto H_i^i\}$$

- ▶ Infinite descending chain:  $\sigma_0 >^{\{F, G\}} \sigma_1 >^{\{F, G\}} \sigma_2 >^{\{F, G\}} \dots$



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .



# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ The problem is of third order.

# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ The problem is of third order.
- ▶ Higher-order unification of the order 3 and above is of type 0.

# Higher Order Unification Is of Type 0

- ▶ Unification problem:  $\Gamma = \{F(\lambda x. G(x), a) \doteq? F(\lambda x. G(x), b)\}$ .
- ▶ The problem is of third order.
- ▶ Higher-order unification of the order 3 and above is of type 0.
- ▶ Second order unification is infinitary.

# Higher Order Unification Is Undecidable

- ▶ Idea: Reduce Hilbert's 10th problem to a higher-order unification problem.
- ▶ Hilbert's 10th problem is undecidable: There is no algorithm that takes as input two polynomials  $P(X_1, \dots, X_n)$  and  $Q(X_1, \dots, X_n)$  with natural coefficients and answers if there exist natural numbers  $m_1, \dots, m_n$  such that

$$P(m_1, \dots, m_n) = Q(m_1, \dots, m_n).$$

- ▶ Reduction requires to represent
  - ▶ natural numbers,
  - ▶ addition,
  - ▶ multiplicationin terms of higher-order unification.



# Higher Order Unification Is Undecidable

Representation (Goldfarb 1981):

- ▶ Natural number  $n$  represented as a  $\lambda$ -term denoted by  $\bar{n}$ :

$$\lambda x.g(a, g(a, \dots g(a, x) \dots))$$

with  $n$  occurrences of  $g$  and  $a$ . The type of  $g$  is  $i \rightarrow i \rightarrow i$  and the type of  $a$  is  $i$ . Such terms are called Goldfarb numbers.

- ▶ Goldfarb numbers are exactly those that solve the unification problem

$$\{g(a, X(a)) \doteq? X(g(a, a))\}$$

and have the type  $i \rightarrow i$ .



# Higher Order Unification Is Undecidable

Representation:

- ▶ Addition is represented by the  $\lambda$ -term *add*:

$$\lambda n.\lambda m.\lambda x. n(m(x)).$$

- ▶ Multiplication is represented by the higher-order unification problem

$$\{Y(a, b, g(g(X_3(a), X_2(b)), a)) \stackrel{?}{=} g(g(a, b), Y(X_1(a), g(a, b), a)) \\ Y(b, a, g(g(X_3(b), X_2(a)), a)) \stackrel{?}{=} g(g(b, a), Y(X_1(b), g(a, a), a))\}$$

that has a solution  $\{X_1 \mapsto \overline{m}_1, X_2 \mapsto \overline{m}_2, X_3 \mapsto \overline{m}_3, Y \mapsto t\}$   
iff  $m_1 \times m_2 = m_3$ .



# Higher Order Unification Is Undecidable

Reduction from Hilbert's 10th problem:

- ▶ Every equation  $P(X_1, \dots, X_n) = Q(X_1, \dots, X_n)$  can be decomposed into a system of equations of the form:

$$X_i + X_j = X_k, \quad X_i \times X_j = X_k, \quad X_i = m.$$

- ▶ With each such system associate a unification problem containing
  - ▶ for each  $X_i$  an equation  $g(a, X_i(a)) \stackrel{?}{=} X_i(g(a, a))$ ,
  - ▶ for each  $X_i + X_j = X_k$  the equation  $add(X_i, X_j) \stackrel{?}{=} X_k$ ,
  - ▶ for each  $X_i \times X_j = X_k$  the two equations used to define multiplication,
  - ▶ for each  $X_i = m$  the equation  $X_i \stackrel{?}{=} \bar{m}$ .



# Second Order Unification Is Undecidable

- ▶ The reduction implies undecidability of higher-order unification.
- ▶ Since the reduction is actually to second-order unification, the result is sharper:

## Theorem

*Second-order unification is undecidable.*

For the details of undecidability of second-order unification, see



[W. D. Goldfarb](#)

The undecidability of the second-order unification problem.

[Theoretical Computer Science 13, 225–230.](#)





# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:

# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:
  1. Enumerate all substitutions (in fact, it is enough to enumerate all closed substitutions).

# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:
  1. Enumerate all substitutions (in fact, it is enough to enumerate all closed substitutions).
  2. For a given unification problem, take the first untried substitution and check whether it is a solution.



# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:
  1. Enumerate all substitutions (in fact, it is enough to enumerate all closed substitutions).
  2. For a given unification problem, take the first untried substitution and check whether it is a solution.
  3. If yes, stop with success. If not, mark the substitution as tried and iterate.



# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:
  1. Enumerate all substitutions (in fact, it is enough to enumerate all closed substitutions).
  2. For a given unification problem, take the first untried substitution and check whether it is a solution.
  3. If yes, stop with success. If not, mark the substitution as tried and iterate.
- ▶ Checking is not hard: Apply the substitution to both sides of each equation, normalize, and compare the normal forms.



# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:
  1. Enumerate all substitutions (in fact, it is enough to enumerate all closed substitutions).
  2. For a given unification problem, take the first untried substitution and check whether it is a solution.
  3. If yes, stop with success. If not, mark the substitution as tried and iterate.
- ▶ Checking is not hard: Apply the substitution to both sides of each equation, normalize, and compare the normal forms.
- ▶ If the problem is solvable, the procedure will detect it after finite steps.



# Higher-Order Unification Procedure

- ▶ Higher-order semi-decision procedure is easy to design:
  1. Enumerate all substitutions (in fact, it is enough to enumerate all closed substitutions).
  2. For a given unification problem, take the first untried substitution and check whether it is a solution.
  3. If yes, stop with success. If not, mark the substitution as tried and iterate.
- ▶ Checking is not hard: Apply the substitution to both sides of each equation, normalize, and compare the normal forms.
- ▶ If the problem is solvable, the procedure will detect it after finite steps.
- ▶ Then... why to bother with looking for another unification procedure?



# Higher-Order Unification Procedure

Why to look for a “better” procedure?



# Higher-Order Unification Procedure

Why to look for a “better” procedure?

- ▶ To find solutions faster.

# Higher-Order Unification Procedure

Why to look for a “better” procedure?

- ▶ To find solutions faster.
- ▶ To report failure for many unsolvable cases.



# Higher-Order Unification Procedure

Why to look for a “better” procedure?

- ▶ To find solutions faster.
- ▶ To report failure for many unsolvable cases.
- ▶ To reduce redundancy.
- ▶ etc.





# Important Observation

- ▶ Flex-flex equation has a form

$$\lambda x_1 \dots \lambda x_k. F(s_1, \dots, s_n) \stackrel{!}{=} \lambda x_1 \dots \lambda x_k. G(t_1, \dots, t_m).$$

The head of both sides are free variables.

# Important Observation

- ▶ Flex-flex equation has a form

$$\lambda x_1 \dots \lambda x_k. F(s_1, \dots, s_n) \stackrel{!}{=} \lambda x_1 \dots \lambda x_k. G(t_1, \dots, t_m).$$

The head of both sides are free variables.

- ▶ Any flex-flex equation is solvable. Just take

$$\{F \mapsto \lambda y_1 \dots \lambda y_n. c, G \mapsto \lambda y_1 \dots \lambda y_m. c\}.$$









# Important Observation

- ▶ Flex-flex equation has a form

$$\lambda x_1 \dots \lambda x_k. F(s_1, \dots, s_n) \stackrel{?}{=} \lambda x_1 \dots \lambda x_k. G(t_1, \dots, t_m).$$

The head of both sides are free variables.

- ▶ Any flex-flex equation is solvable. Just take

$$\{F \mapsto \lambda y_1 \dots \lambda y_n. c, G \mapsto \lambda y_1 \dots \lambda y_m. c\}.$$

- ▶ The appropriate  $c$  always exists because for each type we have at least one constant of that type.
- ▶ Flex-flex equations may introduce infinite branching in the search tree (very undesirable property).
- ▶ Idea: Do not try to solve flex-flex equations. Assume them solved. Preunification.



# Preunification

## Preunifier

- ▶ Let  $\cong$  be the least congruence relation on the set of  $\lambda$ -terms that contains the set of flex-flex pairs.
- ▶ A substitution  $\sigma$  is a preunifier for a unification problem  $\{s_1 \doteq? t_1, \dots, s_n \doteq? t_n\}$  iff

$$\text{normal-form}(s_i\sigma) \cong \text{normal-form}(t_i\sigma)$$

for each  $1 \leq i \leq n$ .

## Convention

- ▶  $\overline{x_n}$  abbreviates  $x_1, \dots, x_n$ .
- ▶  $\lambda\overline{x_n}$  abbreviates  $\lambda x_1 \dots \lambda x_n$ .



# One Technical Notion

## Partial Binding

A partial binding of type  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  ( $U$  atomic) is a term of the form

$$\lambda \overline{x_n}. I(\overline{\lambda y_{m_1}^1} \cdot H_1(\overline{x_n}, \overline{y_{m_1}^1}), \dots, \overline{\lambda y_{m_k}^k} \cdot H_k(\overline{x_n}, \overline{y_{m_k}^k}))$$

where  $I$  is a constant or a variable, and



# One Technical Notion

## Partial Binding

A partial binding of type  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  ( $U$  atomic) is a term of the form

$$\lambda \overline{x_n}. l(\overline{\lambda y_{m_1}^1} \cdot H_1(\overline{x_n}, \overline{y_{m_1}^1}), \dots, \overline{\lambda y_{m_k}^k} \cdot H_k(\overline{x_n}, \overline{y_{m_k}^k}))$$

where  $l$  is a constant or a variable, and

- ▶ the type of  $x_i$  is  $T_i$  for  $1 \leq i \leq n$ ,



# One Technical Notion

## Partial Binding

A partial binding of type  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  ( $U$  atomic) is a term of the form

$$\lambda \overline{x_n}. l(\overline{\lambda y_{m_1}^1} \cdot H_1(\overline{x_n}, \overline{y_{m_1}^1}), \dots, \overline{\lambda y_{m_k}^k} \cdot H_k(\overline{x_n}, \overline{y_{m_k}^k}))$$

where  $l$  is a constant or a variable, and

- ▶ the type of  $x_i$  is  $T_i$  for  $1 \leq i \leq n$ ,
- ▶ the type of  $l$  is  $S_1 \rightarrow \dots \rightarrow S_k \rightarrow U$ , where  $S_j$  is  $R_j^1 \rightarrow \dots \rightarrow R_{m_j}^j \rightarrow S'_j$  ( $S'_j$  atomic) for  $1 \leq j \leq k$ ,



# One Technical Notion

## Partial Binding

A partial binding of type  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  ( $U$  atomic) is a term of the form

$$\lambda \overline{x_n}. l(\overline{\lambda y_{m_1}^1} \cdot H_1(\overline{x_n}, \overline{y_{m_1}^1}), \dots, \overline{\lambda y_{m_k}^k} \cdot H_k(\overline{x_n}, \overline{y_{m_k}^k}))$$

where  $l$  is a constant or a variable, and

- ▶ the type of  $x_i$  is  $T_i$  for  $1 \leq i \leq n$ ,
- ▶ the type of  $l$  is  $S_1 \rightarrow \dots \rightarrow S_k \rightarrow U$ , where  $S_j$  is  $R_j^1 \rightarrow \dots \rightarrow R_{m_j}^j \rightarrow S'_j$  ( $S'_j$  atomic) for  $1 \leq j \leq k$ ,
- ▶ the type of  $y_j^i$  is  $R_j^i$  for  $1 \leq i \leq k$  and  $1 \leq j \leq m_i$ .



# One Technical Notion

## Partial Binding

A partial binding of type  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow U$  ( $U$  atomic) is a term of the form

$$\lambda \overline{x_n}. l(\overline{\lambda y_{m_1}^1} \cdot H_1(\overline{x_n}, \overline{y_{m_1}^1}), \dots, \overline{\lambda y_{m_k}^k} \cdot H_k(\overline{x_n}, \overline{y_{m_k}^k}))$$

where  $l$  is a constant or a variable, and

- ▶ the type of  $x_i$  is  $T_i$  for  $1 \leq i \leq n$ ,
- ▶ the type of  $l$  is  $S_1 \rightarrow \dots \rightarrow S_k \rightarrow U$ , where  $S_i$  is  $R_i^1 \rightarrow \dots \rightarrow R_{m_i}^i \rightarrow S'_i$  ( $S'_i$  atomic) for  $1 \leq i \leq k$ ,
- ▶ the type of  $y_j^i$  is  $R_j^i$  for  $1 \leq i \leq k$  and  $1 \leq j \leq m_i$ .
- ▶ the type of  $H_i$  is  $T_1 \rightarrow \dots \rightarrow T_n \rightarrow R_1^i \rightarrow \dots \rightarrow R_{m_i}^i \rightarrow S'_i$  for  $1 \leq i \leq k$ .



# Partial Binding

$$\lambda \overline{x}_n. I(\lambda \overline{y}_{m_1}^1. H_1(\overline{x}_n, \overline{y}_{m_1}^1), \dots, \lambda \overline{y}_{m_k}^k. H_k(\overline{x}_n, \overline{y}_{m_k}^k))$$

- ▶ Imitation binding:  $I$  is a constant or a free variable.
- ▶ ( $i^{th}$ ) Projection binding:  $I$  is  $x_i$ .
- ▶ A partial binding  $t$  is appropriate to  $F$  if  $t$  and  $F$  have the same types.
- ▶  $F \mapsto t$ : Appropriate partial (imitation, projection) binding if  $t$  is partial (imitation, projection) binding appropriate to  $F$ .





# Higher-Order Preunification Procedure

- ▶ The inference system  $\mathcal{U}_{HOP}$  consists of the rules:
  - ▶ **Trivial**
  - ▶ **Decomposition**
  - ▶ **Variable Elimination**
  - ▶ **Orient**
  - ▶ **Imitation**
  - ▶ **Projection**
- ▶ The rules transform systems: pairs  $\Gamma; \sigma$ , where  $\Gamma$  is a higher-order unification problem and  $\sigma$  is a substitution.
- ▶ A system  $\Gamma; \sigma$  is in presolved form if  $\Gamma$  is either empty or consists of flex-flex equations only.



# Higher-Order Preunification Procedure. Rules

**Trivial:**  $\{t \doteq^? t\} \cup P'; \vartheta \Longrightarrow P'; \vartheta$



# Higher-Order Preunification Procedure. Rules

**Trivial:**  $\{t \doteq^? t\} \cup P'; \vartheta \Longrightarrow P'; \vartheta$

**Decomposition:**

$$\begin{aligned} & \{\lambda \bar{x}_k. l(s_1, \dots, s_n) \doteq^? \lambda \bar{x}_k. l(t_1, \dots, t_n)\} \cup P'; \vartheta \Longrightarrow \\ & \{\lambda \bar{x}_k. s_1 \doteq^? \lambda \bar{x}_k. t_1, \dots, \lambda \bar{x}_k. s_n \doteq^? \lambda \bar{x}_k. t_n\} \cup P'; \vartheta. \end{aligned}$$

where  $l$  is either a constant or one of the bound variables  $x_1, \dots, x_k$ .



# Higher-Order Preunification Procedure. Rules

**Trivial:**  $\{t \doteq^? t\} \cup P'; \vartheta \Longrightarrow P'; \vartheta$

**Decomposition:**

$$\begin{aligned} & \{\lambda \overline{x}_k. l(s_1, \dots, s_n) \doteq^? \lambda \overline{x}_k. l(t_1, \dots, t_n)\} \cup P'; \vartheta \Longrightarrow \\ & \{\lambda \overline{x}_k. s_1 \doteq^? \lambda \overline{x}_k. t_1, \dots, \lambda \overline{x}_k. s_n \doteq^? \lambda \overline{x}_k. t_n\} \cup P'; \vartheta. \end{aligned}$$

where  $l$  is either a constant or one of the bound variables  $x_1, \dots, x_k$ .

**Variable Elimination:**

$$\{\lambda x_1 \dots \lambda x_k. F(x_1, \dots, x_k) \doteq^? t\} \cup P'; \vartheta \Longrightarrow P' \{F \mapsto t\}; \vartheta \{F \mapsto t\}.$$

If  $F \notin fvars(t)$



# Higher-Order Preunification Procedure. Rules

**Orient:**

$$\{\lambda \bar{x}_k. I(t_1, \dots, t_m) \doteq^? \lambda \bar{x}_k. F(s_1, \dots, s_n)\} \cup P'; \vartheta \implies$$
$$\{\lambda \bar{x}_k. F(s_1, \dots, s_n) \doteq^? \lambda \bar{x}_k. I(t_1, \dots, t_m)\} \cup P'; \vartheta$$

where  $I$  is not a free variable.



# Higher-Order Preunification Procedure. Rules

## Orient:

$$\{\lambda \bar{x}_k. I(t_1, \dots, t_m) \doteq^? \lambda \bar{x}_k. F(s_1, \dots, s_n)\} \cup P'; \vartheta \implies$$
$$\{\lambda \bar{x}_k. F(s_1, \dots, s_n) \doteq^? \lambda \bar{x}_k. I(t_1, \dots, t_m)\} \cup P'; \vartheta$$

where  $I$  is not a free variable.

## Imitation:

$$\{\lambda \bar{x}_k. F(s_1, \dots, s_n) \doteq^? \lambda \bar{x}_k. f(t_1, \dots, t_m)\} \cup P'; \vartheta \implies$$
$$\{\lambda \bar{x}_k. f(\lambda \bar{z}_1^1. H_1(s_1, \dots, s_n, \bar{z}_1^1), \dots, \lambda \bar{z}_m^m. H_m(s_1, \dots, s_n, \bar{z}_m^m))\sigma$$
$$\doteq^? \lambda \bar{x}_k. f(t_1, \dots, t_m)\sigma\} \cup P'\sigma; \vartheta\sigma$$

where

- ▶  $\sigma = \{F \mapsto \lambda \bar{y}_n. f(\lambda \bar{z}_1^1. H_1(\bar{y}_n, \bar{z}_1^1), \dots, \lambda \bar{z}_m^m. H_m(\bar{y}_n, \bar{z}_m^m))\}$ , appropriate imitation binding.
- ▶  $H_1, \dots, H_m$  are fresh variables.



# Higher-Order Preunification Procedure. Rules

## Projection:

$$\begin{aligned} & \{\lambda \overline{x}_k. F(s_1, \dots, s_n) \doteq^? \lambda \overline{x}_k. l(t_1, \dots, t_m)\} \cup P'; \vartheta \implies \\ & \{\lambda \overline{x}_k. s_i(\lambda \overline{z}_{r_1}^1. H_1(s_1, \dots, s_n, \overline{z}_{r_1}^1), \dots, \lambda \overline{z}_{r_m}^m. H_m(s_1, \dots, s_n, \overline{z}_{r_m}^m))\sigma \\ & \quad \doteq^? \lambda \overline{x}_k. l(t_1, \dots, t_m)\sigma\} \cup P'\sigma; \vartheta\sigma \end{aligned}$$

where

- ▶  $l$  is either a constant or one of the bound variables  $x_1, \dots, x_k$ ,
- ▶  $\sigma = \{F \mapsto \lambda \overline{y}_n. y_i(\lambda \overline{z}_{r_1}^1. H_1(\overline{y}_n, \overline{z}_{r_1}^1), \dots, \lambda \overline{z}_{r_m}^m. H_m(\overline{y}_n, \overline{z}_{r_m}^m))\}$ , appropriate projection binding.
- ▶  $H_1, \dots, H_m$  are fresh variables.



# Higher-Order Preunification Procedure. Control

In order to solve a higher-order unification problem  $\Gamma$ :

- ▶ Create an initial system  $\Gamma; \varepsilon$ .





# Higher-Order Preunification Procedure. Control

In order to solve a higher-order unification problem  $\Gamma$ :

- ▶ Create an initial system  $\Gamma; \varepsilon$ .
- ▶ Apply successively rules from  $\mathcal{U}_{HOP}$ , building a complete (finitely branching, but potentially infinite) tree of derivations.



# Higher-Order Preunification Procedure. Control

In order to solve a higher-order unification problem  $\Gamma$ :

- ▶ Create an initial system  $\Gamma; \varepsilon$ .
- ▶ Apply successively rules from  $\mathcal{U}_{HOP}$ , building a complete (finitely branching, but potentially infinite) tree of derivations.
- ▶ If no rule can be applied to a node, and it contains at least one equation that is not flex-flex, then extend the branch with  $\perp$ , indicating failure.



# Higher-Order Preunification Procedure. Control

In order to solve a higher-order unification problem  $\Gamma$ :

- ▶ Create an initial system  $\Gamma; \varepsilon$ .
- ▶ Apply successively rules from  $\mathcal{U}_{HOP}$ , building a complete (finitely branching, but potentially infinite) tree of derivations.
- ▶ If no rule can be applied to a node, and it contains at least one equation that is not flex-flex, then extend the branch with  $\perp$ , indicating failure.
- ▶ Successful leaves contain presolved systems.



# Higher-Order Preunification Procedure. Control

In order to solve a higher-order unification problem  $\Gamma$ :

- ▶ Create an initial system  $\Gamma; \varepsilon$ .
- ▶ Apply successively rules from  $\mathcal{U}_{HOP}$ , building a complete (finitely branching, but potentially infinite) tree of derivations.
- ▶ If no rule can be applied to a node, and it contains at least one equation that is not flex-flex, then extend the branch with  $\perp$ , indicating failure.
- ▶ Successful leaves contain presolved systems.
- ▶ If  $\Delta; \sigma$  is a successful leaf,  $\sigma$  is a solution of  $\Gamma$  computed by the higher-order preunification procedure.



# Higher-Order Preunification. Major Results

## Theorem (Soundness)

*If  $\Gamma; \varepsilon \Longrightarrow^* \Delta; \sigma$  and  $\Delta$  is in presolved form, then  $\sigma|_{fvars(\Gamma)}$  is a preunifier of  $\Gamma$ .*

## Theorem (Completeness)

*If  $\vartheta$  is a preunifier of  $\Gamma$ , then there exists a sequence of transformations  $\Gamma; \varepsilon \Longrightarrow^* \Delta; \sigma$  such that  $\Delta$  is in presolved form, and  $\sigma \leq_{\beta}^{fvars(\Gamma)} \vartheta$ .*



# Higher-Order Preunification. Optimization

- ▶ The procedure can be optimized by stripping off the binder  $\lambda x$  when  $x$  does not occur in the body.
- ▶ For instance, Elimination rule does not apply to  $\lambda x.\lambda y. P(x) \doteq? \lambda x.\lambda y. f(a)$
- ▶ After removing  $\lambda y$  from both sides, Elimination can be applied directly.



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\{F(f(a)) \doteq^? f(F(a))\}; \varepsilon$$





# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\begin{aligned} & \{F(f(a)) \doteq? f(F(a))\}; \varepsilon \\ & \implies_{Proj} \{f(a) \doteq? f(a)\}; \{F \mapsto \lambda x. x\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\begin{aligned} & \{F(f(a)) \doteq^? f(F(a))\}; \varepsilon \\ & \implies_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. x\} \\ & \implies_{Tr} \emptyset; \{F \mapsto \lambda x. x\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\begin{aligned} & \{F(f(a)) \doteq^? f(F(a))\}; \varepsilon \\ & \implies_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. x\} \\ & \implies_{Tr} \emptyset; \{F \mapsto \lambda x. x\} \end{aligned}$$

$$\{F(f(a)) \doteq^? f(F(a))\}; \varepsilon$$

# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\begin{aligned} & \{F(f(a)) \doteq^? f(F(a))\}; \varepsilon \\ & \implies_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. x\} \\ & \implies_{Tr} \emptyset; \{F \mapsto \lambda x. x\} \end{aligned}$$

$$\begin{aligned} & \{F(f(a)) \doteq^? f(F(a))\}; \varepsilon \\ & \implies_{lmit} \{f(G(f(a))) \doteq^? f(f(G(a)))\}; \{F \mapsto \lambda x. f(G(x))\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\begin{aligned} & \{F(f(a)) \doteq^? f(F(a))\}; \varepsilon \\ & \implies_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. x\} \\ & \implies_{Tr} \emptyset; \{F \mapsto \lambda x. x\} \end{aligned}$$

$$\begin{aligned} & \{F(f(a)) \doteq^? f(F(a))\}; \varepsilon \\ & \implies_{Limit} \{f(G(f(a))) \doteq^? f(f(G(a)))\}; \{F \mapsto \lambda x. f(G(x))\} \\ & \implies_{Dec} \{G(f(a)) \doteq^? f(G(a))\}; \{F \mapsto \lambda x. f(G(x))\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\{F(f(a)) \doteq^? f(F(a))\}; \varepsilon$$

$$\Longrightarrow_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. x\}$$

$$\Longrightarrow_{Tr} \emptyset; \{F \mapsto \lambda x. x\}$$

$$\{F(f(a)) \doteq^? f(F(a))\}; \varepsilon$$

$$\Longrightarrow_{Limit} \{f(G(f(a))) \doteq^? f(f(G(a)))\}; \{F \mapsto \lambda x. f(G(x))\}$$

$$\Longrightarrow_{Dec} \{G(f(a)) \doteq^? f(G(a))\}; \{F \mapsto \lambda x. f(G(x))\}$$

$$\Longrightarrow_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. f(x), G \mapsto \lambda x. x\}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Unification problem  $\{F(f(a)) \doteq^? f(F(a))\}$ .
- ▶ The preunification procedure enumerates the complete set of (pre)unifiers that is infinite.
- ▶ Here we show only two derivations.

$$\{F(f(a)) \doteq^? f(F(a))\}; \varepsilon$$

$$\Longrightarrow_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. x\}$$

$$\Longrightarrow_{Tr} \emptyset; \{F \mapsto \lambda x. x\}$$

$$\{F(f(a)) \doteq^? f(F(a))\}; \varepsilon$$

$$\Longrightarrow_{Limit} \{f(G(f(a))) \doteq^? f(f(G(a)))\}; \{F \mapsto \lambda x. f(G(x))\}$$

$$\Longrightarrow_{Dec} \{G(f(a)) \doteq^? f(G(a))\}; \{F \mapsto \lambda x. f(G(x))\}$$

$$\Longrightarrow_{Proj} \{f(a) \doteq^? f(a)\}; \{F \mapsto \lambda x. f(x), G \mapsto \lambda x. x\}$$

$$\Longrightarrow_{Tr} \emptyset; \{F \mapsto \lambda x. f(x), G \mapsto \lambda x. x\}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}$ .
- ▶ Here we show only the successful derivation.



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}$ .
- ▶ Here we show only the successful derivation.

$$\{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}; \varepsilon$$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}$ .
- ▶ Here we show only the successful derivation.

$$\begin{aligned} & \{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}; \varepsilon \\ & \implies_{\text{limit}} \{\lambda x. g(H_1(f(x, G)), H_2(f(x, G))) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}; \\ & \quad \{F \mapsto \lambda y. g(H_1(y), H_2(y))\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}$ .
- ▶ Here we show only the successful derivation.

$$\{\lambda x. F(f(x, G)) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}; \varepsilon$$

$$\begin{aligned} \implies_{\text{Imit}} & \{\lambda x. g(H_1(f(x, G)), H_2(f(x, G))) \doteq? \lambda x. g(f(x, G_1), f(x, G_2))\}; \\ & \{F \mapsto \lambda y. g(H_1(y), H_2(y))\} \end{aligned}$$

$$\begin{aligned} \implies_{\text{Dec, Proj, Proj}} & \{\lambda x. f(x, G) \doteq? \lambda x. f(x, G_1), \lambda x. f(x, G) \doteq? \lambda x. f(x, G_2)\}; \\ & \{F \mapsto \lambda y. g(y, y), H_1 \mapsto \lambda y. y, H_2 \mapsto \lambda y. y\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(f(x, G)) \doteq^? \lambda x. g(f(x, G_1), f(x, G_2))\}$ .
- ▶ Here we show only the successful derivation.

$$\{\lambda x. F(f(x, G)) \doteq^? \lambda x. g(f(x, G_1), f(x, G_2))\}; \varepsilon$$

$$\begin{aligned} \implies_{\text{Imit}} & \{\lambda x. g(H_1(f(x, G)), H_2(f(x, G))) \doteq^? \lambda x. g(f(x, G_1), f(x, G_2))\}; \\ & \{F \mapsto \lambda y. g(H_1(y), H_2(y))\} \end{aligned}$$

$$\begin{aligned} \implies_{\text{Dec, Proj, Proj}} & \{\lambda x. f(x, G) \doteq^? \lambda x. f(x, G_1), \lambda x. f(x, G) \doteq^? \lambda x. f(x, G_2)\}; \\ & \{F \mapsto \lambda y. g(y, y), H_1 \mapsto \lambda y. y, H_2 \mapsto \lambda y. y\} \end{aligned}$$

$$\begin{aligned} \implies_{\text{Dec, Tr, Dec, Tr}} & \{\lambda x. G \doteq^? \lambda x. G_1, \lambda x. G \doteq^? \lambda x. G_2\}; \\ & \{F \mapsto \lambda y. g(y, y), H_1 \mapsto \lambda y. y, H_2 \mapsto \lambda y. y\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(f(x, G)) \doteq^? \lambda x. g(f(x, G_1), f(x, G_2))\}$ .
- ▶ Here we show only the successful derivation.

$$\begin{aligned} & \{\lambda x. F(f(x, G)) \doteq^? \lambda x. g(f(x, G_1), f(x, G_2))\}; \varepsilon \\ \implies & \text{Imit} \{\lambda x. g(H_1(f(x, G)), H_2(f(x, G))) \doteq^? \lambda x. g(f(x, G_1), f(x, G_2))\}; \\ & \{F \mapsto \lambda y. g(H_1(y), H_2(y))\} \\ \implies & \text{Dec, Proj, Proj} \{\lambda x. f(x, G) \doteq^? \lambda x. f(x, G_1), \lambda x. f(x, G) \doteq^? \lambda x. f(x, G_2)\}; \\ & \{F \mapsto \lambda y. g(y, y), H_1 \mapsto \lambda y. y, H_2 \mapsto \lambda y. y\} \\ \implies & \text{Dec, Tr, Dec, Tr} \{\lambda x. G \doteq^? \lambda x. G_1, \lambda x. G \doteq^? \lambda x. G_2\}; \\ & \{F \mapsto \lambda y. g(y, y), H_1 \mapsto \lambda y. y, H_2 \mapsto \lambda y. y\} \\ \implies & \text{Elim}^2 \emptyset; \{F \mapsto \lambda y. g(y, y), H_1 \mapsto \lambda y. y, H_2 \mapsto \lambda y. y, G \mapsto G_2, G_1 \mapsto G_2\} \end{aligned}$$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}$ .
- ▶ One of the successful derivations.



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}$ .
- ▶ One of the successful derivations.

$\{\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}; \varepsilon$



# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}$ .
- ▶ One of the successful derivations.

$\{\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}; \varepsilon$

$\implies_{limit} \{\lambda x. f(H(x, a)) \doteq? \lambda x. f(G(a, x))\}; \{F \mapsto \lambda y_1. \lambda y_2. f(H(y_1, y_2))\}$





# Higher-Order Preunification. Examples

## Example

- ▶ Problem  $\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}$ .
- ▶ One of the successful derivations.

$\{\{\lambda x. F(x, a) \doteq? \lambda x. f(G(a, x))\}; \varepsilon$

$\implies_{\text{Imit}} \{\lambda x. f(H(x, a)) \doteq? \lambda x. f(G(a, x))\}; \{F \mapsto \lambda y_1. \lambda y_2. f(H(y_1, y_2))\}$

$\implies_{\text{Dec}} \{\lambda x. H(x, a) \doteq? \lambda x. G(a, x)\}; \{F \mapsto \lambda y_1. \lambda y_2. f(H(y_1, y_2))\}$

Flex-flex.



# Decidable Subcases

Some decidable subcases of higher-order unification:

- ▶ Monadic second-order unification. Terms do not contain constants of arity greater than 1.  
Example:  $\{\lambda x.f(F(x)) \doteq? \lambda x.F(f(x))\}$ .
- ▶ Second-order unification with linear occurrences of second-order variables.
- ▶ Unification with higher-order patterns. Pattern is a term  $t$  such that for every subterm of the form  $F(s_1, \dots, s_n)$ , the  $s$ 's are distinct variables bound in  $t$ .  
Example:  $\{\lambda x.\lambda y.F(x) \doteq? \lambda x.\lambda y.c(G(y, x))\}$ .
- ▶ Higher-order matching. One side in the equations is a closed term.  
Example.  $\{\lambda x.F(x, \lambda y.y) \doteq? \lambda x.f(x, a)\}$ .
- ▶ Stratified second-order unification.
- ▶ Bounded second-order unification.

