

Loops and Legs in QFT 2014, Weimar, Germany

Recent Symbolic Summation Methods to Solve Coupled Systems of Differential and Difference Equations

Carsten Schneider

RISC, J. Kepler University Linz, Austria

joint work with A. Behring, J. Blümlein, A. De Freitas (DESY, Zeuthen)

April 29, 2014

Recent symbolic summation methods

A challenging diagram

A new method for coupled systems

A general tactic

Feynman integrals

A general tactic

Feynman integrals

↓ non-trivial transformations (DESY)

multiple sums

A general tactic

Feynman integrals

↓ non-trivial transformations (DESY)

multiple sums

↓ symbolic summation

compact expression in terms
of special functions

A general tactic

Feynman integrals

↓ non-trivial transformations (DESY)

multiple sums

↓ symbolic summation

compact expression in terms
of special functions

Tactic 1: Expand the summand and simplify

Tactic 1: Expand and simplify

$$\text{GIVEN } F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! \times$$

$$\times \underbrace{B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right)}_{f(N, k)} \binom{N}{k}$$

where

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

Tactic 1: Expand and simplify

$$\text{GIVEN } F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! \times$$

$$\underbrace{\times B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}}_{f(N, k)}$$

FIND the first coefficients of the ε -expansion

$$F(N) \stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \dots$$

Tactic 1: Expand and simplify

$$\text{GIVEN } F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! \times$$

$$\underbrace{\times B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}}_{f(N, k)}$$

Step 1: Compute the first coefficients of the ε -expansion

$$f(N, k) = f_{-3}(N, k) \varepsilon^{-3} + f_{-2}(N, k) \varepsilon^{-2} + f_{-1}(N, k) \varepsilon^{-1} +$$

Tactic 1: Expand and simplify

$$\text{GIVEN } F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! \times$$

$$\underbrace{\times B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}}_{f(N, k)}$$

Step 2: **Simplify** the sums in

$$\sum_{k=1}^N f(N, k) = \left(\sum_{k=1}^{\infty} f_{-3}(N, k)\right) \varepsilon^{-3} + \left(\sum_{k=1}^{\infty} f_{-2}(N, k)\right) \varepsilon^{-2} + \left(\sum_{k=1}^{\infty} f_{-1}(N, k)\right) \varepsilon^{-1} +$$

Simplify

$$F_{-1}(N) = \sum_{k=1}^N (-1)^{k+1} \binom{N}{k} \left(\frac{(2+3k)(-2+3k+7k^2+3k^3)}{3k^2(1+k)^3} + \frac{2S_2(k)}{1+k} + \frac{\zeta_2}{2(1+k)} \right)$$

where

$$S_a(N) = \sum_{i=1}^N \frac{\text{sign}(a)^i}{i^a} \quad \text{and} \quad \zeta_a = \sum_{i=1}^{\infty} \frac{1}{i^a}$$

Simplify

$$F_{-1}(N) = \sum_{k=1}^N (-1)^{k+1} \binom{N}{k} \left(\frac{(2+3k)(-2+3k+7k^2+3k^3)}{3k^2(1+k)^3} + \frac{2S_2(k)}{1+k} + \frac{\zeta_2}{2(1+k)} \right)$$

↓ (summation package Sigma.m)

$$\begin{aligned} & (16N^3 + 144N^2 + 413N + 384)(N+1)^2 F_{-1}(N) \\ & - (N+2)(2N+5)(16N^3 + 112N^2 + 221N + 113) F_{-1}(N+1) \\ & + (N+3)^2(16N^3 + 96N^2 + 173N + 99) F_{-1}(N+2) \\ & = \frac{1}{2}(4N^2 + 21N + 29)\zeta_2 + \frac{-64N^5 - 500N^4 - 1133N^3 + 203N^2 + 3516N + 3090}{3(N+2)(N+3)} \end{aligned}$$

Simplify

$$F_{-1}(N) = \sum_{k=1}^N (-1)^{k+1} \binom{N}{k} \left(\frac{(2+3k)(-2+3k+7k^2+3k^3)}{3k^2(1+k)^3} + \frac{2S_2(k)}{1+k} + \frac{\zeta_2}{2(1+k)} \right)$$

↓ (summation package Sigma.m)

$$\begin{aligned} & (16N^3 + 144N^2 + 413N + 384)(N+1)^2 F_{-1}(N) \\ & - (N+2)(2N+5)(16N^3 + 112N^2 + 221N + 113) F_{-1}(N+1) \\ & + (N+3)^2(16N^3 + 96N^2 + 173N + 99) F_{-1}(N+2) \\ & = \frac{1}{2}(4N^2 + 21N + 29)\zeta_2 + \frac{-64N^5 - 500N^4 - 1133N^3 + 203N^2 + 3516N + 3090}{3(N+2)(N+3)} \end{aligned}$$

↓ (summation package Sigma.m)

$$\begin{aligned} & \left\{ c_1 \frac{1-4N}{N+1} + c_2 \frac{-14N-13}{(N+1)^2} \right. \\ & + \frac{(4N-1)S_1(N)}{N+1} + \frac{(1-4N)S_1(N)^2}{6(N+1)} + \frac{(14N+13)S_1(N)}{3(N+1)^2} \\ & \left. + \frac{175N^2 + 334N + 155}{12(N+1)^3} + \frac{(1-4N)S_2(N)}{6(N+1)} + \frac{\zeta_2}{8(N+1)} \mid c_1, c_2 \in \mathbb{Q} \right\} \end{aligned}$$

Simplify

$$F_{-1}(N) = \sum_{k=1}^N (-1)^{k+1} \binom{N}{k} \left(\frac{(2+3k)(-2+3k+7k^2+3k^3)}{3k^2(1+k)^3} + \frac{2S_2(k)}{1+k} + \frac{\zeta_2}{2(1+k)} \right)$$



$$\left\{ c_1 \frac{1-4N}{N+1} + c_2 \frac{-14N-13}{(N+1)^2} + \frac{(4N-1)S_1(N)}{N+1} + \frac{(1-4N)S_1(N)^2}{6(N+1)} + \frac{(14N+13)S_1(N)}{3(N+1)^2} + \frac{175N^2+334N+155}{12(N+1)^3} + \frac{(1-4N)S_2(N)}{6(N+1)} + \frac{\zeta_2}{8(N+1)} \mid c_1, c_2 \in \mathbb{Q} \right\}$$

Simplify

$$F_{-1}(N) = \sum_{k=1}^N (-1)^{k+1} \binom{N}{k} \left(\frac{(2+3k)(-2+3k+7k^2+3k^3)}{3k^2(1+k)^3} + \frac{2S_2(k)}{1+k} + \frac{\zeta_2}{2(1+k)} \right)$$

||

$$\begin{aligned} & \left(\frac{1}{12} - \frac{1}{8}\zeta_2 \right) \frac{1-4N}{N+1} + 1 \frac{-14N-13}{(N+1)^2} \\ & + \frac{(4N-1)S_1(N)}{N+1} + \frac{(1-4N)S_1(N)^2}{6(N+1)} + \frac{(14N+13)S_1(N)}{3(N+1)^2} \\ & + \frac{175N^2+334N+155}{12(N+1)^3} + \frac{(1-4N)S_2(N)}{6(N+1)} + \frac{\zeta_2}{8(N+1)} \end{aligned}$$

1. Creative telescoping (for the special case of hypergeometric terms see Zeilberger's algorithm (1991))

GIVEN a definite sum

$$F(N) = \sum_{k=0}^N f(N, k);$$

$f(N, k)$: indefinite nested product-sum in k ;
 N : extra parameter

FIND a recurrence for $F(N)$

1. Creative telescoping (for the special case of hypergeometric terms see Zeilberger's algorithm (1991))

GIVEN a **definite** sum

$$F(N) = \sum_{k=0}^N f(N, k);$$

$f(N, k)$: indefinite nested product-sum in k ;
 N : extra parameter

FIND a **recurrence** for $F(N)$

2. Recurrence solving

GIVEN a recurrence

$a_0(N), \dots, a_d(N), h(N)$:
indefinite nested product-sum expressions.

$$a_0(N)F(N) + \dots + a_d(N)F(N + d) = h(N);$$

FIND **all solutions** expressible by indefinite nested products/sums

(Abramov/Bronstein/Petkovšek/CS, in preparation)

1. Creative telescoping (for the special case of hypergeometric terms see Zeilberger's algorithm (1991))

GIVEN a definite sum

$$F(N) = \sum_{k=0}^N f(N, k);$$

$f(N, k)$: indefinite nested product-sum in k ;
 N : extra parameter

FIND a recurrence for $F(N)$

2. Recurrence solving

GIVEN a recurrence

$a_0(N), \dots, a_d(N), h(N)$:
 indefinite nested product-sum expressions.


$$a_0(N)F(N) + \dots + a_d(N)F(N + d) = h(N);$$

FIND all solutions expressible by indefinite nested products/sums
 (Abramov/Bronstein/Petkovšek/CS, in preparation)

3. Find a "closed form"

$F(N)$ =combined solutions in terms of indefinite nested sums.

Consider a massive 3-loop ladder graph (Ablinger, Blümlein, Hasselhuhn, Klein, CS, Wissbrock, 2012)



$$= F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \boxed{F_0(N)}$$

Consider a massive 3-loop ladder graph (Ablinger, Blümlein, Hasselhuhn, Klein, CS, Wissbrock, 2012)



$$= F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \boxed{F_0(N)}$$

Simplify

||

$$\sum_{j=0}^{N-3} \sum_{k=0}^j \sum_{l=0}^k \sum_{q=0}^{-j+N-3} \sum_{s=1}^{-l+N-q-3} \sum_{r=0}^{-l+N-q-s-3} (-1)^{-j+k-l+N-q-3} \times$$

$$\times \frac{\binom{j+1}{k+1} \binom{k}{l} \binom{N-1}{j+2} \binom{-j+N-3}{q} \binom{-l+N-q-3}{s} \binom{-l+N-q-s-3}{r} r! (-l+N-q-r-s-3)! (s-1)!}{(-l+N-q-2)! (-j+N-1) (N-q-r-s-2) (q+s+1)}$$

$$\left[4S_1(-j+N-1) - 4S_1(-j+N-2) - 2S_1(k) \right.$$

$$\left. - (S_1(-l+N-q-2) + S_1(-l+N-q-r-s-3) - 2S_1(r+s)) \right.$$

$$\left. + 2S_1(s-1) - 2S_1(r+s) \right] + \mathbf{3 \text{ further 6-fold sums}}$$

$$\boxed{F_0(N)} = \quad (\text{using Sigma.m, EvaluateMultiSums.m and J. Ablinger's HarmonicSums.m package})$$

$$\begin{aligned}
 & \frac{7}{12}S_1(N)^4 + \frac{(17N+5)S_1(N)^3}{3N(N+1)} + \left(\frac{35N^2-2N-5}{2N^2(N+1)^2} + \frac{13S_2(N)}{2} + \frac{5(-1)^N}{2N^2} \right) S_1(N)^2 \\
 & + \left(-\frac{4(13N+5)}{N^2(N+1)^2} + \left(\frac{4(-1)^N(2N+1)}{N(N+1)} - \frac{13}{N} \right) S_2(N) + \left(\frac{29}{3} - (-1)^N \right) S_3(N) \right. \\
 & + \left(2 + 2(-1)^N \right) S_{2,1}(N) - 28S_{-2,1}(N) + \frac{20(-1)^N}{N^2(N+1)} S_1(N) + \left(\frac{3}{4} + (-1)^N \right) S_2(N)^2 \\
 & - 2(-1)^N S_{-2}(N)^2 + S_{-3}(N) \left(\frac{2(3N-5)}{N(N+1)} + (26 + 4(-1)^N) S_1(N) + \frac{4(-1)^N}{N+1} \right) \\
 & + \left(\frac{(-1)^N(5-3N)}{2N^2(N+1)} - \frac{5}{2N^2} \right) S_2(N) + S_{-2}(N) (10S_1(N)^2 + \left(\frac{8(-1)^N(2N+1)}{N(N+1)} \right. \\
 & + \left. \frac{4(3N-1)}{N(N+1)} \right) S_1(N) + \frac{8(-1)^N(3N+1)}{N(N+1)^2} + (-22 + 6(-1)^N) S_2(N) - \frac{16}{N(N+1)} \\
 & + \left(\frac{(-1)^N(9N+5)}{N(N+1)} - \frac{29}{3N} \right) S_3(N) + \left(\frac{19}{2} - 2(-1)^N \right) S_4(N) + (-6 + 5(-1)^N) S_{-4}(N) \\
 & + \left(-\frac{2(-1)^N(9N+5)}{N(N+1)} - \frac{2}{N} \right) S_{2,1}(N) + (20 + 2(-1)^N) S_{2,-2}(N) + (-17 + 13(-1)^N) S_{3,1}(N) \\
 & - \frac{8(-1)^N(2N+1) + 4(9N+1)}{N(N+1)} S_{-2,1}(N) - (24 + 4(-1)^N) S_{-3,1}(N) + (3 - 5(-1)^N) S_{2,1,1}(N) \\
 & + 32S_{-2,1,1}(N) + \left(\frac{3}{2} S_1(N)^2 - \frac{3S_1(N)}{N} + \frac{3}{2} (-1)^N S_{-2}(N) \right) \zeta_2
 \end{aligned}$$

A general tactic

Feynman integrals

↓ non-trivial transformations (DESY)

multiple sums

↓ symbolic summation

compact expression in terms
of special functions

Tactic 2: Expand a recurrence in ε

$$F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}$$

↓ (summation package Sigma.m)

$$2(N+1)^2 F(N) + (3\varepsilon^2 + 3\varepsilon N + 9\varepsilon - 4N^2 - 12N - 8) F(N+1) - (2\varepsilon - N - 1)(\varepsilon + 2N + 6) F(N+2) = 0\varepsilon^{-3} - \frac{16}{3}\varepsilon^{-2} + \frac{40}{3}\varepsilon^{-1} - \left(2\zeta_2 - \frac{68}{3}\right)\varepsilon^0 + \dots$$

$$F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}$$

↓ (summation package Sigma.m)

$$2(N+1)^2 F(N) + (3\varepsilon^2 + 3\varepsilon N + 9\varepsilon - 4N^2 - 12N - 8) F(N+1) - (2\varepsilon - N - 1)(\varepsilon + 2N + 6) F(N+2) = 0\varepsilon^{-3} - \frac{16}{3}\varepsilon^{-2} + \frac{40}{3}\varepsilon^{-1} - \left(2\zeta_2 - \frac{68}{3}\right)\varepsilon^0 + \dots$$

$$F(1) = \frac{2}{3}\varepsilon^{-3} - \frac{11}{6}\varepsilon^{-2} + \left(\frac{\zeta_2}{4} + \frac{79}{24}\right)\varepsilon^{-1} + \dots$$

$$F(2) = \frac{8}{9}\varepsilon^{-3} - \frac{73}{27}\varepsilon^{-2} + \left(\frac{\zeta_2}{3} + \frac{1415}{324}\right)\varepsilon^{-1} + \dots$$

Ansatz (for power series)

$$\begin{aligned}
 & a_0(\varepsilon, N) \left[F_0(N) + F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\
 & + a_1(\varepsilon, N) \left[F_0(N+1) + F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\
 & + \\
 & \vdots \\
 & + a_d(\varepsilon, N) \left[F_0(N+d) + F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\
 & \qquad \qquad \qquad = h_0(N) + h_1(N)\varepsilon + h_1(N)\varepsilon^2 + \dots
 \end{aligned}$$

given (in terms of indefinite nested sums and products)

Ansatz (for power series)

$$\begin{aligned} & a_0(\varepsilon, N) \left[F_0(N) + F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\ & + a_1(\varepsilon, N) \left[F_0(N+1) + F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\ & + \\ & \vdots \\ & + a_d(\varepsilon, N) \left[F_0(N+d) + F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\ & \qquad \qquad \qquad = h_0(N) + h_1(N)\varepsilon + h_1(N)\varepsilon^2 + \dots \end{aligned}$$

↓ constant terms must agree

$$a_0(0, N)F_0(N) + a_1(0, N)F_0(N+1) + \dots + a_d(0, N)F_0(N+d) = h_0(N)$$

Ansatz (for power series)

$$\begin{aligned}
 & a_0(\varepsilon, N) \left[F_0(N) + F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\
 & + a_1(\varepsilon, N) \left[F_0(N+1) + F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\
 & + \\
 & \vdots \\
 & + a_d(\varepsilon, N) \left[F_0(N+d) + F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\
 & \qquad \qquad \qquad = h_0(N) + h_1(N)\varepsilon + h_1(N)\varepsilon^2 + \dots
 \end{aligned}$$

↓ constant terms must agree

$$a_0(0, N)F_0(N) + a_1(0, N)F_0(N+1) + \dots + a_d(0, N)F_0(N+d) = h_0(N)$$

REC solver: Using the initial values $F_0(1), F_0(2), \dots$ determine $F_0(N)$ in terms of indefinite nested sums and products.

Ansatz (for power series)

$$\begin{aligned}
 & a_0(\varepsilon, N) \left[F_0(N) + F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\
 & + a_1(\varepsilon, N) \left[F_0(N+1) + F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\
 & + \\
 & \vdots \\
 & + a_d(\varepsilon, N) \left[F_0(N+d) + F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\
 & \qquad \qquad \qquad = h_0(N) + h_1(N)\varepsilon + h_1(N)\varepsilon^2 + \dots
 \end{aligned}$$

↓ constant terms must agree

$$a_0(0, N)F_0(N) + a_1(0, N)F_0(N+1) + \dots + a_d(0, N)F_0(N+d) = h_0(N)$$

Ansatz (for power series)

$$\begin{aligned}
 & a_0(\varepsilon, N) \left[F_0(N) + F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\
 & + a_1(\varepsilon, N) \left[F_0(N+1) + F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\
 & + \\
 & \vdots \\
 & + a_d(\varepsilon, N) \left[F_0(N+d) + F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\
 & \qquad \qquad \qquad = h_0(N) + h_1(N)\varepsilon + h_1(N)\varepsilon^2 + \dots
 \end{aligned}$$

↓ constant terms must agree

$$a_0(0, N)F_0(N) + a_1(0, N)F_0(N+1) + \dots + a_d(0, N)F_0(N+d) = h_0(N)$$

$$\begin{aligned} & a_0(\varepsilon, N) \left[F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\ & + a_1(\varepsilon, N) \left[F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\ & + \\ & \vdots \\ & + a_d(\varepsilon, N) \left[F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\ & \qquad \qquad \qquad = h'_0(N) + h'_1(N)\varepsilon + h'_2(N)\varepsilon^2 + \dots \end{aligned}$$

$$\begin{aligned}
 & a_0(\varepsilon, N) \left[F_1(N)\varepsilon + F_2(N)\varepsilon^2 + \dots \right] \\
 & + a_1(\varepsilon, N) \left[F_1(N+1)\varepsilon + F_2(N+1)\varepsilon^2 + \dots \right] \\
 & + \\
 & \vdots \\
 & + a_d(\varepsilon, N) \left[F_1(N+d)\varepsilon + F_2(N+d)\varepsilon^2 + \dots \right] \\
 & \qquad \qquad \qquad = \underbrace{h'_0(N)}_{=0} + h'_1(N)\varepsilon + h'_2(N)\varepsilon^2 + \dots
 \end{aligned}$$

Devide by ε

$$\begin{aligned} & a_0(\varepsilon, N) \left[F_1(N) + F_2(N)\varepsilon + \dots \right] \\ & + a_1(\varepsilon, N) \left[F_1(N+1) + F_2(N+1)\varepsilon + \dots \right] \\ & + \\ & \vdots \\ & + a_d(\varepsilon, N) \left[F_1(N+d) + F_2(N+d)\varepsilon + \dots \right] = h'_1(N) + h'_2(N)\varepsilon + \dots \end{aligned}$$

Now repeat for $F_1(N), F_2(N), \dots$

Remark: Works the same for Laurent series.

[Blümlein, Klein, CS, Stan, 2012; arXiv:1011.2656v2]

$$F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}$$

↓ (summation package Sigma.m)

$$2(N+1)^2 F(N) + (3\varepsilon^2 + 3\varepsilon N + 9\varepsilon - 4N^2 - 12N - 8) F(N+1) - (2\varepsilon - N - 1)(\varepsilon + 2N + 6) F(N+2) = 0\varepsilon^{-3} - \frac{16}{3}\varepsilon^{-2} + \frac{40}{3}\varepsilon^{-1} - \left(2\zeta_2 - \frac{68}{3}\right)\varepsilon^0 + \dots$$

$$F(1) = \frac{2}{3}\varepsilon^{-3} - \frac{11}{6}\varepsilon^{-2} + \left(\frac{\zeta_2}{4} + \frac{79}{24}\right)\varepsilon^{-1} + \dots$$

$$F(2) = \frac{8}{9}\varepsilon^{-3} - \frac{73}{27}\varepsilon^{-2} + \left(\frac{\zeta_2}{3} + \frac{1415}{324}\right)\varepsilon^{-1} + \dots$$

$$F(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}$$

↓ (summation package Sigma.m)

$$2(N+1)^2 F(N) + (3\varepsilon^2 + 3\varepsilon N + 9\varepsilon - 4N^2 - 12N - 8) F(N+1) - (2\varepsilon - N - 1)(\varepsilon + 2N + 6) F(N+2) = 0\varepsilon^{-3} - \frac{16}{3}\varepsilon^{-2} + \frac{40}{3}\varepsilon^{-1} - (2\zeta_2 - \frac{68}{3})\varepsilon^0 + \dots$$

$$F(1) = \frac{2}{3}\varepsilon^{-3} - \frac{11}{6}\varepsilon^{-2} + \left(\frac{\zeta_2}{4} + \frac{79}{24}\right)\varepsilon^{-1} + \dots$$

$$F(2) = \frac{8}{9}\varepsilon^{-3} - \frac{73}{27}\varepsilon^{-2} + \left(\frac{\zeta_2}{3} + \frac{1415}{324}\right)\varepsilon^{-1} + \dots$$

↓ (summation package Sigma.m)

$$F(N) = \frac{4N}{3(N+1)}\varepsilon^{-3} - \left(\frac{2(2N+1)}{3(N+1)}S_1(N) + \frac{2N(2N+3)}{3(N+1)^2}\right)\varepsilon^{-2}$$

$$\left(\frac{(1-4N)}{6(N+1)}S_1(N)^2 - \frac{N(N^2-2)}{3(N+1)^3} + \frac{(3N+2)(4N+5)}{3(N+1)^2}S_1(N) + \frac{(1-4N)}{6(N+1)}S_2(N) + \frac{N\zeta_2}{2(N+1)}\right)\varepsilon^{-1} + \dots$$

Find a recurrence for the integral/sum

$$F(N) = \int_0^1 \dots \int_0^1 \Phi(\varepsilon, N, x_1, x_2, \dots, x_7) dx_1 dx_2 \dots dx_7$$
$$\stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \dots$$

multivariate
Almquist/Zeilberger
(Jakob Ablinger)

$$a_0(\varepsilon, N)F(N) + \dots + a_d(\varepsilon, N)F(N + d) = h(\varepsilon, N)$$

Find a recurrence for the integral/sum

$$F(N) = \int_0^1 \dots \int_0^1 \Phi(\varepsilon, N, x_1, x_2, \dots, x_7) dx_1 dx_2 \dots dx_7$$

$$\stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \dots$$

multivariate
Almquist/Zeilberger
(Jakob Ablinger)

$$\sum_{i_1} \dots \sum_{i_7} f(\varepsilon, N, i_1, i_2, \dots, i_7)$$

MultiSum Package
(Flavia Stan)

$$a_0(\varepsilon, N)F(N) + \dots + a_d(\varepsilon, N)F(N + d) = h(\varepsilon, N)$$

Find a recurrence for the integral/sum

$$F(N) = \int_0^1 \dots \int_0^1 \Phi(\varepsilon, N, x_1, x_2, \dots, x_7) dx_1 dx_2 \dots dx_7$$

$$\stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \dots$$

multivariate
Almquist/Zeilberger
(Jakob Ablinger)

$$\sum_{i_1} \dots \sum_{i_7} f(\varepsilon, N, i_1, i_2, \dots, i_7)$$

MultiSum Package
(Flavia Stan)

Holonomic/difference field Approach
(Mark Round)

$$a_0(\varepsilon, N)F(N) + \dots + a_d(\varepsilon, N)F(N + d) = h(\varepsilon, N)$$

Find a recurrence for the integral/sum

$$F(N) = \int_0^1 \dots \int_0^1 \Phi(\varepsilon, N, x_1, x_2, \dots, x_7) dx_1 dx_2 \dots dx_7$$

$$\stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \dots$$

 ε -recurrence solver

multivariate
Almquist/Zeilberger
(Jakob Ablinger)

$$\sum_{i_1} \dots \sum_{i_7} f(\varepsilon, N, i_1, i_2, \dots, i_7)$$


MultiSum Package
(Flavia Stan)

Holonomic/difference field Approach
(Mark Round)

$$a_0(\varepsilon, N)F(N) + \dots + a_d(\varepsilon, N)F(N + d) = h(\varepsilon, N)$$

A challenging diagram and an algorithm for coupled systems

A challenging diagram (ladder graph with 6 massive fermion lines)




$$\stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + F_0(N)\varepsilon^0 + \dots$$

Strategies:

- Symbolic summation tools: failed (so far) 😞

A challenging diagram (ladder graph with 6 massive fermion lines)



$$\stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + F_0(N)\varepsilon^0 + \dots$$


Strategies:

- ▶ Symbolic summation tools: failed (so far) ☹️
- ▶ Brown's hyperlogarithm algorithm: works for the scalar version where

$$\lim_{\varepsilon \rightarrow 0} D_4(N) = F_0(N).$$

[Ablinger, Blümlein, Raab, Schneider, Wissbrock, 2014; arXiv:1403.1137 [hep-ph]]

A challenging diagram (ladder graph with 6 massive fermion lines)



$$D_4(N) \stackrel{?}{=} F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + F_0(N)\varepsilon^0 + \dots$$

Strategies:

- ▶ Symbolic summation tools: failed (so far) ☹️
- ▶ Brown's hyperlogarithm algorithm: works for the scalar version where

$$\lim_{\varepsilon \rightarrow 0} D_4(N) = F_0(N).$$

[Ablinger, Blümlein, Raab, Schneider, Wissbrock, 2014; arXiv:1403.1137 [hep-ph]]

- ▶ New approach: for the complete diagram

Consider the power series of $D_4(N)$:

$$D_4(N) \longrightarrow \hat{D}_4(x) = \sum_{N=0}^{\infty} D_4(N)x^N$$

(holonomic closure properties)

Consider the power series of $D_4(N)$:

$$D_4(N) \longleftarrow \hat{D}_4(x) = \sum_{N=0}^{\infty} D_4(N)x^N$$

(holonomic closure properties)

IBP (extension of REDUZE_2, A.v. Manteuffel) gives

$$\underbrace{\sum_{N=0}^{\infty} D_4(N)x^N}_{\hat{D}_4(x)} = \frac{(1545842\varepsilon^5 x^5 - 14325922\varepsilon^5 x^4 \dots + 1524096x^2 - 653184x)}{23328\varepsilon^2(x-1)x^5} \hat{B}_1(x) + \square \hat{B}_2(x) + \dots + \square \hat{B}_{52}(x) \dots$$

$\hat{B}_1(x), \dots, \hat{B}_{52}(x)$ can be handled with sophisticated Mellin-Barnes techniques (DESY colleagues) and symbolic summation (see first slides).

IBP (extension of REDUZE_2, A.v. Manteuffel) gives

$$\underbrace{\sum_{N=0}^{\infty} D_4(N)x^N}_{\hat{D}_4(x)} = \frac{(1545842\varepsilon^5 x^5 - 14325922\varepsilon^5 x^4 \dots + 1524096x^2 - 653184x)}{23328\varepsilon^2(x-1)x^5} \hat{B}_1(x) + \square \hat{B}_2(x) + \dots + \square \hat{B}_{52}(x)$$

$\hat{B}_1(x), \dots, \hat{B}_{52}(x)$ can be handled with sophisticated Mellin-Barnes techniques (DESY colleagues) and symbolic summation (see first slides).

E.g.,

$$\hat{B}_1(x) = \sum_{N=0}^{\infty} B_1(N)x^N$$

with

$$B_1(N) = \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k}$$

IBP (extension of REDUZE_2, A.v. Manteuffel) gives

$$\underbrace{\sum_{N=0}^{\infty} D_4(N)x^N}_{\hat{D}_4(x)} = \boxed{\frac{(1545842\varepsilon^5 x^5 - 14325922\varepsilon^5 x^4 \dots + 1524096x^2 - 653184x)}{23328\varepsilon^2(x-1)x^5}} \hat{B}_1(x) + \square \hat{B}_2(x) + \dots + \square \hat{B}_{52}(x)$$

$\hat{B}_1(x), \dots, \hat{B}_{52}(x)$ can be handled with sophisticated Mellin-Barnes techniques (DESY colleagues) and symbolic summation (see first slides).

E.g.,

$$\hat{B}_1(x) = \sum_{N=0}^{\infty} B_1(N)x^N$$

with

$$\begin{aligned} B_1(N) &= \sum_{k=1}^N (-1)^k e^{-\frac{3\varepsilon\gamma}{2}} \left(-2 - \frac{3\varepsilon}{2}\right)! B\left(2+k, \frac{\varepsilon}{2}\right) B(-\varepsilon+k, -\varepsilon) B\left(1 - \frac{\varepsilon}{2} + k, 1 + \frac{\varepsilon}{2}\right) \binom{N}{k} \\ &= \frac{4N}{3(N+1)} \varepsilon^{-3} - \left(\frac{2(2N+1)}{3(N+1)} S_1(N) + \frac{2N(2N+3)}{3(N+1)^2}\right) \varepsilon^{-2} + \dots \end{aligned}$$

IBP (extension of REDUZE_2, A.v. Manteuffel) gives

$$\underbrace{\sum_{N=0}^{\infty} D_4(N)x^N}_{\hat{D}_4(x)} = \boxed{\frac{(1545842\epsilon^5 x^5 - 14325922\epsilon^5 x^4 + \dots + 1524096x^2 - 653184x)}{23328\epsilon^2(x-1)x^5}} \hat{B}_1(x) \\
 + \square \hat{B}_2(x) + \dots + \square \hat{B}_{52}(x) \\
 + \boxed{-\frac{(122\epsilon^4 x^3 - 2647\epsilon^4 x^2 + \dots - 304\epsilon + 24x^3 - 24x)}{4\epsilon x^4}} \hat{I}_1(x) \\
 + \boxed{\frac{(589\epsilon^5 x^3 - 20123\epsilon^5 x^2 + \dots - 896\epsilon + 96x^3 - 96x)}{16\epsilon^2 x^4}} \hat{I}_2(x) \\
 + \boxed{\frac{(589\epsilon^5 x^3 - 21509\epsilon^5 x^2 + \dots - 1152\epsilon + 96x^3 - 96x)}{16\epsilon^2 x^4}} \hat{I}_3(x) \\
 + \square \hat{I}_4(x) + \dots + \square \hat{I}_{15}(x)$$

However, $\hat{I}_1(x), \dots, \hat{I}_{15}(x)$ are hard to handle. Luckily...

... there are differential relations among the integrals. E.g.,

$$D_x \hat{I}_1(x) = - \frac{(-\varepsilon+x-1)}{(x-1)x} \hat{I}_1(x) - \frac{2}{(x-1)x} \hat{I}_2(x) \\ + \frac{1}{(x-1)x} \hat{B}_1(x) + \dots$$

$$D_x \hat{I}_2(x) = \frac{(3(\varepsilon+4)^2 - 22(\varepsilon+4) + 40)}{4(x-1)} \hat{I}_1(x) \\ + \frac{(-(\varepsilon+4)(3x-1) + 9x-2)}{2(x-1)x} \hat{I}_2(x) - \frac{(\varepsilon+1)}{2(x-1)} \hat{I}_3(x) \\ + \frac{1}{4} \frac{((\varepsilon+4)^2(6x-25) - 2(\varepsilon+4)(17x-75) + 48x-224)}{(5\varepsilon+6)(x-1)x} \hat{B}_1(x) + \dots$$

$$D_x \hat{I}_3(x) = - \frac{(3(\varepsilon+4)^2(x-2) - 22(\varepsilon+4)(x-2) + 40x-80)}{4(x-1)x} \hat{I}_1(x) \\ + \frac{((\varepsilon+4)(3x-5) - 11x+18)}{2(x-1)x} \hat{I}_2(x) - \frac{(-(\varepsilon+4)(x-2) + 5x-8)}{2(x-1)x} \hat{I}_3(x) \\ - \frac{1}{4} \frac{((\varepsilon+4)^2(6x-25) - 2(\varepsilon+4)(17x-75) + 48x-224)}{(5\varepsilon+6)(x-1)x} \hat{B}_1(x) + \dots$$

... there are differential relations among the integrals. E.g.,

$$D_x \hat{I}_1(x) = - \frac{(-\varepsilon+x-1)}{(x-1)x} \hat{I}_1(x) - \frac{2}{(x-1)x} \hat{I}_2(x) \\ + \frac{1}{(x-1)x} \hat{B}_1(x) + \dots$$

$$D_x \hat{I}_2(x) = \frac{(3(\varepsilon+4)^2 - 22(\varepsilon+4) + 40)}{4(x-1)} \hat{I}_1(x) \\ + \frac{(-(\varepsilon+4)(3x-1) + 9x-2)}{2(x-1)x} \hat{I}_2(x) - \frac{(\varepsilon+1)}{2(x-1)} \hat{I}_3(x) \\ + \frac{1}{4} \frac{((\varepsilon+4)^2(6x-25) - 2(\varepsilon+4)(17x-75) + 48x-224)}{(5\varepsilon+6)(x-1)x} \hat{B}_1(x) + \dots$$

$$D_x \hat{I}_3(x) = - \frac{(3(\varepsilon+4)^2(x-2) - 22(\varepsilon+4)(x-2) + 40x-80)}{4(x-1)x} \hat{I}_1(x) \\ + \frac{((\varepsilon+4)(3x-5) - 11x+18)}{2(x-1)x} \hat{I}_2(x) - \frac{(-(\varepsilon+4)(x-2) + 5x-8)}{2(x-1)x} \hat{I}_3(x) \\ - \frac{1}{4} \frac{((\varepsilon+4)^2(6x-25) - 2(\varepsilon+4)(17x-75) + 48x-224)}{(5\varepsilon+6)(x-1)x} \hat{B}_1(x) + \dots$$

Step 1: From a DE system to a REC system

$$\begin{aligned}D_x \hat{I}_1(x) &= - \frac{(-\varepsilon+x-1)}{(x-1)x} \hat{I}_1(x) \\ &\quad - \frac{2}{(x-1)x} \hat{I}_2(x) \\ &\quad + \frac{1}{(x-1)x} \hat{B}_1(x) + \dots\end{aligned}$$

Step 1: From a DE system to a REC system

$$\begin{aligned} D_x \sum_{N=0}^{\infty} I_1(N)x^N &= - \frac{(-\varepsilon+x-1)}{(x-1)x} \sum_{N=0}^{\infty} I_1(N)x^N \\ &\quad - \frac{2}{(x-1)x} \sum_{N=0}^{\infty} I_2(N)x^N \\ &\quad + \frac{1}{(x-1)x} \sum_{N=0}^{\infty} B_1(N)x^N + \dots \end{aligned}$$

Step 1: From a DE system to a REC system

$$\begin{aligned} \sum_{N=1}^{\infty} I_1(N) N x^{N-1} &= - \frac{(-\varepsilon+x-1)}{(x-1)x} \sum_{N=0}^{\infty} I_1(N) x^N \\ &\quad - \frac{2}{(x-1)x} \sum_{N=0}^{\infty} I_2(N) x^N \\ &\quad + \frac{1}{(x-1)x} \sum_{N=0}^{\infty} B_1(N) x^N + \dots \end{aligned}$$

Step 1: From a DE system to a REC system

$$\begin{aligned} \sum_{N=1}^{\infty} I_1(N) N x^{N-1} &= - \frac{(-\varepsilon+x-1)}{(x-1)x} \sum_{N=0}^{\infty} I_1(N) x^N \\ &\quad - \frac{2}{(x-1)x} \sum_{N=0}^{\infty} I_2(N) x^N \\ &\quad + \frac{1}{(x-1)x} \sum_{N=0}^{\infty} B_1(N) x^N + \dots \end{aligned}$$

↓ N th coefficient

$$N I_1(N-1) - (\varepsilon + N + 1) I_1(N) + 2 I_2(N) = B_1(N) + \dots$$

... there are differential relations among the integrals. E.g.,

$$NI_1(N-1) - (\varepsilon + N + 1)I_1(N) + 2I_2(N) \\ = B_1(N) + \dots$$

$$D_x \hat{I}_2(x) = \frac{(3(\varepsilon+4)^2 - 22(\varepsilon+4) + 40)}{4(x-1)} \hat{I}_1(x) \\ + \frac{(-(\varepsilon+4)(3x-1) + 9x-2)}{2(x-1)x} \hat{I}_2(x) - \frac{(\varepsilon+1)}{2(x-1)} \hat{I}_3(x) \\ + \frac{1}{4} \frac{((\varepsilon+4)^2(6x-25) - 2(\varepsilon+4)(17x-75) + 48x-224)}{(5\varepsilon+6)(x-1)x} \hat{B}_1(x) + \dots$$

$$D_x \hat{I}_3(x) = - \frac{(3(\varepsilon+4)^2(x-2) - 22(\varepsilon+4)(x-2) + 40x-80)}{4(x-1)x} \hat{I}_1(x) \\ + \frac{((\varepsilon+4)(3x-5) - 11x+18)}{2(x-1)x} \hat{I}_2(x) - \frac{(-(\varepsilon+4)(x-2) + 5x-8)}{2(x-1)x} \hat{I}_3(x) \\ - \frac{1}{4} \frac{((\varepsilon+4)^2(6x-25) - 2(\varepsilon+4)(17x-75) + 48x-224)}{(5\varepsilon+6)(x-1)x} \hat{B}_1(x) + \dots$$

A coupled system of difference equations

$$NI_1(N-1) - (\varepsilon + N + 1)I_1(N) + 2I_2(N) \\ = B_1(N) + \dots$$

$$2(\varepsilon + 2N + 2)I_2(N) - 2(3\varepsilon + 2N + 1)I_2(N-1) \\ + \varepsilon(3\varepsilon + 2)I_1(N-1) - 2(\varepsilon + 1)I_3(N-1) \\ = (5\varepsilon + 4)B_1(N) - \frac{2(\varepsilon + 1)(3\varepsilon + 4)}{5\varepsilon + 6}B_1(N-1) + \dots$$

$$4(\varepsilon - N)I_3(N) - 2\varepsilon(3\varepsilon + 2)I_1(N) + \varepsilon(3\varepsilon + 2)I_1(N-1) \\ - 2(3\varepsilon + 1)I_2(N-1) + 2(5\varepsilon + 2)I_2(N) \\ - 2(\varepsilon - 2N + 1)I_3(N-1) \\ = -\frac{2(\varepsilon + 1)(3\varepsilon + 4)}{5\varepsilon + 6}B_1(N-1) + (5\varepsilon + 4)B_1(N) + \dots$$

A coupled system of difference equations

$$\begin{aligned}
 & NI_1(N-1) - (\varepsilon + N + 1)I_1(N) + 2I_2(N) \\
 &= + \frac{4(N+2)}{3(N+1)}\varepsilon^{-3} + \left(\frac{2(2N+1)}{3(N+1)}S_1(N) - \frac{2(6N^2+13N+8)}{3(N+1)^2} \right)\varepsilon^{-2} + \dots
 \end{aligned}$$

$$\begin{aligned}
 & 2(\varepsilon + 2N + 2)I_2(N) - 2(3\varepsilon + 2N + 1)I_2(N-1) \\
 & \quad + \varepsilon(3\varepsilon + 2)I_1(N-1) - 2(\varepsilon + 1)I_3(N-1) \\
 &= \frac{8}{3}\varepsilon^{-3} + \left(\frac{4}{3}S_1(N)^2 - \frac{4(N+1)}{N}S_1(N) + \frac{4}{3}S_2(N) + \zeta_2 + 6 \right)\varepsilon^{-1} + \dots
 \end{aligned}$$

$$\begin{aligned}
 & 4(\varepsilon - N)I_3(N) - 2\varepsilon(3\varepsilon + 2)I_1(N) + \varepsilon(3\varepsilon + 2)I_1(N-1) \\
 & \quad - 2(3\varepsilon + 1)I_2(N-1) + 2(5\varepsilon + 2)I_2(N) \\
 & \quad - 2(\varepsilon - 2N + 1)I_3(N-1) \\
 &= - \frac{8}{3}\varepsilon^{-3} - \left(\frac{8}{3}S_1(N) - 4 \right)\varepsilon^{-2} \\
 & \quad - \left(\frac{4}{3}S_1(N)^2 - \frac{4(N+1)}{N}S_1(N) + \frac{4}{3}S_2(N) + \zeta_2 + 6 \right)\varepsilon^{-1} + \dots
 \end{aligned}$$

Step 2: Uncouple the system

$$\square I_1(N-1) + \square I_1(N) + \square I_2(N)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-2} + \square \varepsilon^{-1} + \dots$$

$$\square I_2(N) + \square I_2(N-1) + \square I_1(N-1) + \square I_3(N-1)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-1} + \dots$$

$$\square I_3(N) + \square I_1(N) + \square I_1(N-1) + \square I_2(N) + \square I_3(N-1)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-2} + \square \varepsilon^{-1} + \dots$$

Step 2: Uncouple the system

$$\square I_1(N-1) + \square I_1(N) + \square I_2(N)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-2} + \square \varepsilon^{-1} + \dots$$

$$\square I_2(N) + \square I_2(N-1) + \square I_1(N-1) + \square I_3(N-1)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-1} + \dots$$

$$\square I_3(N) + \square I_1(N) + \square I_1(N-1) + \square I_2(N) + \square I_3(N-1)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-2} + \square \varepsilon^{-1} + \dots$$

↓ (uncoupling algorithms^a, S. Gerhold's OrseSys.m)

$$\square I_1(N) + \square I_1(N+1) + \square I_1(N+2) + \square I_1(N+3)$$

$$= \square \varepsilon^{-3} + \square \varepsilon^{-2} + \square \varepsilon^{-1} + \dots$$

$$I_2(N) = \text{expression in } I_1(N)$$

$$I_3(N) = \text{expression in } I_1(N)$$

^a We use Zürcher's uncoupling algorithm (1994)

More precisely, we get:

$$\begin{aligned} & -2(N+1)(N+2)(\varepsilon+N+2)I_1(N) - (N+2)(2\varepsilon^2 - 5\varepsilon N - 7\varepsilon - 6N^2 - 28N - 32) \\ & + (\varepsilon^3 + 4\varepsilon^2 N + 14\varepsilon^2 - 4\varepsilon N^2 - 13\varepsilon N - 3\varepsilon - 6N^3 - 50N^2 - 136N - 120)I_1(N+2) \\ & - (\varepsilon - N - 2)(\varepsilon + N + 4)(\varepsilon + 2N + 8)I_1(N+3) \\ & = -\frac{4(N+2)}{3(N+3)}\varepsilon^{-3} + \frac{2(4N^4 + 35N^3 + 101N^2 + 105N + 25)}{3(N+1)(N+2)(N+3)^2}\varepsilon^{-2} + \dots \end{aligned}$$

Step 3: Solve the scalar recurrence

$$\begin{aligned}
 & -2(N+1)(N+2)(\varepsilon+N+2)I_1(N) - (N+2)(2\varepsilon^2 - 5\varepsilon N - 7\varepsilon - 6N^2 - 28N - 32) \\
 & + (\varepsilon^3 + 4\varepsilon^2 N + 14\varepsilon^2 - 4\varepsilon N^2 - 13\varepsilon N - 3\varepsilon - 6N^3 - 50N^2 - 136N - 120)I_1(N+2) \\
 & - (\varepsilon - N - 2)(\varepsilon + N + 4)(\varepsilon + 2N + 8)I_1(N+3) \\
 & = -\frac{4(N+2)}{3(N+3)}\varepsilon^{-3} + \frac{2(4N^4 + 35N^3 + 101N^2 + 105N + 25)}{3(N+1)(N+2)(N+3)^2}\varepsilon^{-2} + \dots
 \end{aligned}$$

$$\begin{aligned}
 I_1(1) &= \frac{5}{\varepsilon^3} - \frac{163}{12\varepsilon^2} + \left(\frac{15\zeta_2}{8} + \frac{1223}{48}\right)\varepsilon^{-1} + \dots && \text{using, e.g., an extension of} \\
 I_1(2) &= \frac{130}{27\varepsilon^3} - \frac{695}{54\varepsilon^2} + \left(\frac{65\zeta_2}{36} + \frac{46379}{1944}\right)\varepsilon^{-1} + \dots && \text{MATAD (M. Steinhauser);} \\
 I_1(3) &= \frac{169}{36\varepsilon^3} - \frac{395}{32\varepsilon^2} + \left(\frac{169\zeta_2}{96} + \frac{470071}{20736}\right)\varepsilon^{-1} + \dots && \text{see also A. De Freitas' talk}
 \end{aligned}$$

Step 3: Solve the scalar recurrence

$$\begin{aligned}
 & -2(N+1)(N+2)(\varepsilon+N+2)I_1(N) - (N+2)(2\varepsilon^2 - 5\varepsilon N - 7\varepsilon - 6N^2 - 28N - 32) \\
 & + (\varepsilon^3 + 4\varepsilon^2 N + 14\varepsilon^2 - 4\varepsilon N^2 - 13\varepsilon N - 3\varepsilon - 6N^3 - 50N^2 - 136N - 120)I_1(N+2) \\
 & - (\varepsilon - N - 2)(\varepsilon + N + 4)(\varepsilon + 2N + 8)I_1(N+3) \\
 & = -\frac{4(N+2)}{3(N+3)}\varepsilon^{-3} + \frac{2(4N^4 + 35N^3 + 101N^2 + 105N + 25)}{3(N+1)(N+2)(N+3)^2}\varepsilon^{-2} + \dots
 \end{aligned}$$

$$I_1(1) = \frac{5}{\varepsilon^3} - \frac{163}{12\varepsilon^2} + \left(\frac{15\zeta_2}{8} + \frac{1223}{48}\right)\varepsilon^{-1} + \dots \quad \text{using, e.g., an extension of}$$

$$I_1(2) = \frac{130}{27\varepsilon^3} - \frac{695}{54\varepsilon^2} + \left(\frac{65\zeta_2}{36} + \frac{46379}{1944}\right)\varepsilon^{-1} + \dots \quad \text{MATAD (M. Steinhauser);}$$

$$I_1(3) = \frac{169}{36\varepsilon^3} - \frac{395}{32\varepsilon^2} + \left(\frac{169\zeta_2}{96} + \frac{470071}{20736}\right)\varepsilon^{-1} + \dots \quad \text{see also A. De Freitas' talk}$$

↓ (Sigma.m's recurrence solver, see first slides)

$$\begin{aligned}
 I_1(N) &= \left(\frac{4(3N^2+6N+4)}{3(N+1)^2} + \frac{4S_1(N)}{3(N+1)}\right)\varepsilon^{-3} \\
 &- \left(\frac{2(20N^3+58N^2+57N+22)}{3(N+1)^3} + \frac{S_1(N)^2}{N+1} + \frac{2(N+2)(2N-1)S_1(N)}{3(N+1)^2} - \frac{S_2(N)}{N+1}\right)\varepsilon^{-2} + \dots
 \end{aligned}$$

Step 4: Compute $I_2(N)$ and $I_3(N)$:

Recall: by uncoupling we expressed $I_2(N)$ and $I_3(N)$ by $I_1(N)$

Step 4: Compute $I_2(N)$ and $I_3(N)$:

Recall: by uncoupling we expressed $I_2(N)$ and $I_3(N)$ by $I_1(N)$, i.e.,

$$I_2(N) = \square I_1(N) + \square I_1(N+1) + \square I_1(N+2) \\ - \frac{2(N+2)}{3(N+1)} \varepsilon^{-3} + \left(\frac{6N^3+25N^2+33N+15}{3(N+1)^2(N+2)} + \frac{(-2N-1)}{3(N+1)} S_1(N) \right) \varepsilon^{-2} + \dots$$

$$I_3(N) = \square I_1(N) + \square I_1(N+1) + \square I_1(N+2) \\ + \frac{2(N+2)}{3(N+1)} \varepsilon^{-3} + \left(\frac{-2N^3-3N^2+3N+3}{3(N+1)^2(N+2)} + \frac{(2N+1)}{3(N+1)} S_1(N) \right) \varepsilon^{-2} + \dots$$

Step 4: Compute $I_2(N)$ and $I_3(N)$:

Recall: by uncoupling we expressed $I_2(N)$ and $I_3(N)$ by $I_1(N)$, i.e.,

$$I_2(N) = \square I_1(N) + \square I_1(N+1) + \square I_1(N+2) \\ - \frac{2(N+2)}{3(N+1)} \varepsilon^{-3} + \left(\frac{6N^3+25N^2+33N+15}{3(N+1)^2(N+2)} + \frac{(-2N-1)}{3(N+1)} S_1(N) \right) \varepsilon^{-2} + \dots$$

$$I_3(N) = \square I_1(N) + \square I_1(N+1) + \square I_1(N+2) \\ + \frac{2(N+2)}{3(N+1)} \varepsilon^{-3} + \left(\frac{-2N^3-3N^2+3N+3}{3(N+1)^2(N+2)} + \frac{(2N+1)}{3(N+1)} S_1(N) \right) \varepsilon^{-2} + \dots$$

This yields

$$I_2(N) = \frac{4}{3\varepsilon^3} - \frac{2}{\varepsilon^2} + \left(-\frac{1}{3} S_1(N)^2 + \frac{2}{3} S_1(N) - \frac{1}{3} S_2(N) + \frac{5N+7}{3(N+1)} + \frac{\zeta_2}{2} \right) \varepsilon^{-1} + \dots$$

$$I_3(N) = \frac{8}{3\varepsilon^3} + \left(\frac{4(N+2)}{3(N+1)} S_1(N) - \frac{4(4N^2+7N+2)}{3(N+1)^2} \right) \varepsilon^{-2} \\ + \left(-\frac{2(4N^2+11N+10)}{3(N+1)^2} S_1(N) + \frac{2(12N^3+32N^2+25N+2)}{3(N+1)^3} \right. \\ \left. + \frac{(N-2)}{3(N+1)} S_1(N)^2 + \frac{(N-2)}{3(N+1)} S_2(N) + \zeta_2 \right) \varepsilon^{-1} + \dots$$

Compute the remaining integrals

$$\underbrace{\sum_{N=0}^{\infty} D_4(N)x^N}_{\hat{D}_4(x)} = \boxed{\frac{(1545842\epsilon^5 x^5 - 14325922\epsilon^5 x^4 + \dots + 1524096x^2 - 653184x)}{23328\epsilon^2(x-1)x^5}} \hat{B}_1(x)$$

$$+ \square \hat{B}_2(x) + \dots + \square \hat{B}_{52}(x)$$

$$+ \boxed{-\frac{(122\epsilon^4 x^3 - 2647\epsilon^4 x^2 + \dots - 304\epsilon + 24x^3 - 24x)}{4\epsilon x^4}} \hat{I}_1(x)$$

$$+ \boxed{\frac{(589\epsilon^5 x^3 - 20123\epsilon^5 x^2 + \dots - 896\epsilon + 96x^3 - 96x)}{16\epsilon^2 x^4}} \hat{I}_2(x)$$

$$+ \boxed{\frac{(589\epsilon^5 x^3 - 21509\epsilon^5 x^2 + \dots - 1152\epsilon + 96x^3 - 96x)}{16\epsilon^2 x^4}} \hat{I}_3(x)$$

$$+ \square \hat{I}_4(x) + \dots + \square \hat{I}_{15}(x)$$

Analogously, all $\hat{I}_j(x) = \sum_{N=0}^{\infty} I_j(N)x^N$, $j = 1, \dots, 15$ can be computed.

Final step: Insert all subresults

$$\underbrace{\sum_{N=0}^{\infty} D_4(N)x^N}_{\hat{D}_4(x)} = \boxed{\frac{(1545842\epsilon^5 x^5 - 14325922\epsilon^5 x^4 + \dots + 1524096x^2 - 653184x)}{23328\epsilon^2(x-1)x^5}} \hat{B}_1(x)$$

$$+ \square \hat{B}_2(x) + \dots + \square \hat{B}_{52}(x)$$

$$+ \boxed{-\frac{(122\epsilon^4 x^3 - 2647\epsilon^4 x^2 + \dots - 304\epsilon + 24x^3 - 24x)}{4\epsilon x^4}} \hat{I}_1(x)$$

$$+ \boxed{\frac{(589\epsilon^5 x^3 - 20123\epsilon^5 x^2 + \dots - 896\epsilon + 96x^3 - 96x)}{16\epsilon^2 x^4}} \hat{I}_2(x)$$

$$+ \boxed{\frac{(589\epsilon^5 x^3 - 21509\epsilon^5 x^2 + \dots - 1152\epsilon + 96x^3 - 96x)}{16\epsilon^2 x^4}} \hat{I}_3(x)$$

$$+ \square \hat{I}_4(x) + \dots + \square \hat{I}_{15}(x)$$

Plugging in all expansion and extracting the N -th coefficient
 (using `HarmonicSums.m`, `Sigma.m`, `EvaluateMultiSum.m`, `SumProduction.m`)
 yield

$$I_4(N) = \left(\frac{64(N^2+N-1)}{3(N+1)(N+2)(N+3)(N+4)} - \frac{64S_1(N)}{3(N+3)(N+4)} \right) \mathcal{E}^{-3}$$

$$\begin{aligned}
I_4(N) &= \left(\frac{64(N^2+N-1)}{3(N+1)(N+2)(N+3)(N+4)} - \frac{64S_1(N)}{3(N+3)(N+4)} \right) \mathcal{E}^{-3} \\
&+ \left(\frac{4(5N+27)}{3(N+2)(N+3)(N+4)} S_1(N)^2 - \frac{4(3N^5+68N^4+379N^3+648N^2-98N-696)}{3(N+1)(N+2)^2(N+3)^2(N+4)^2} S_1(N) \right. \\
&+ \left. \frac{4(14N^6+214N^5+1179N^4+3050N^3+4097N^2+3094N+1200)}{3(N+1)^2(N+2)^2(N+3)^2(N+4)^2} + \frac{4(N+1)(4N+17)S_2(N)}{3(N+2)(N+3)(N+4)} \right) \mathcal{E}^{-2}
\end{aligned}$$

$$\begin{aligned}
I_4(N) = & \left(\frac{64(N^2+N-1)}{3(N+1)(N+2)(N+3)(N+4)} - \frac{64S_1(N)}{3(N+3)(N+4)} \right) \mathcal{E}^{-3} \\
& + \left(\frac{4(5N+27)}{3(N+2)(N+3)(N+4)} S_1(N)^2 - \frac{4(3N^5+68N^4+379N^3+648N^2-98N-696)}{3(N+1)(N+2)^2(N+3)^2(N+4)^2} S_1(N) \right. \\
& + \left. \frac{4(14N^6+214N^5+1179N^4+3050N^3+4097N^2+3094N+1200)}{3(N+1)^2(N+2)^2(N+3)^2(N+4)^2} + \frac{4(N+1)(4N+17)S_2(N)}{3(N+2)(N+3)(N+4)} \right) \mathcal{E}^{-2} \\
& + \left(\frac{-23N^2-35N-176}{9(N+2)(N+3)(N+4)(N+5)} S_1(N)^3 - \frac{2(10N^2+53N+106)}{3(N+2)(N+3)(N+4)} S_{2,1}(N) + \zeta_2 \left(\frac{8(N^2+N-1)}{(N+1)(N+2)(N+3)(N+4)} \right) \right. \\
& - \left. \frac{8}{(N+3)(N+4)} S_1(N) \right) + \frac{-8N^3-95N^2-171N-56}{3(N+2)(N+3)(N+4)(N+5)} S_2(N) S_1(N) + \frac{2(30N^3+469N^2+1873N+2542)}{9(N+2)(N+3)(N+4)(N+5)} S_3(N) \\
& + \frac{25N^6+213N^5+491N^4-1007N^3-7942N^2-15988N-10340}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_1(N)^2 \\
& + \frac{-85N^6-1469N^5-8965N^4-23889N^3-25644N^2-3724N+5780}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_2(N) \\
& - \frac{2(94N^{10}+2202N^9+22629N^8+133916N^7+505769N^6+\dots+1817100N+563760)}{3(N+1)^2(N+2)^3(N+3)^3(N+4)^3(N+5)} S_1(N) \\
& - \frac{2(44N^{11}+1696N^{10}+26555N^9+230482N^8+\dots+4371092N+623040)}{3(N+1)^3(N+2)^3(N+3)^3(N+4)^3(N+5)} \Big) \mathcal{E}^{-1}
\end{aligned}$$

$$\begin{aligned}
I_4(N) &= \left(\frac{64(N^2+N-1)}{3(N+1)(N+2)(N+3)(N+4)} - \frac{64S_1(N)}{3(N+3)(N+4)} \right) \varepsilon^{-3} \\
&+ \left(\frac{4(5N+27)}{3(N+2)(N+3)(N+4)} S_1(N)^2 - \frac{4(3N^5+68N^4+379N^3+648N^2-98N-696)}{3(N+1)(N+2)^2(N+3)^2(N+4)^2} S_1(N) \right. \\
&+ \left. \frac{4(14N^6+214N^5+1179N^4+3050N^3+4097N^2+3094N+1200)}{3(N+1)^2(N+2)^2(N+3)^2(N+4)^2} + \frac{4(N+1)(4N+17)S_2(N)}{3(N+2)(N+3)(N+4)} \right) \varepsilon^{-2} \\
&+ \left(\frac{-23N^2-35N-176}{9(N+2)(N+3)(N+4)(N+5)} S_1(N)^3 - \frac{2(10N^2+53N+106)}{3(N+2)(N+3)(N+4)} S_{2,1}(N) + \zeta_2 \left(\frac{8(N^2+N-1)}{(N+1)(N+2)(N+3)(N+4)} \right) \right. \\
&- \frac{8}{(N+3)(N+4)} S_1(N) \left. + \frac{-8N^3-95N^2-171N-56}{3(N+2)(N+3)(N+4)(N+5)} S_2(N) S_1(N) + \frac{2(30N^3+469N^2+1873N+2542)}{9(N+2)(N+3)(N+4)(N+5)} S_3(N) \right. \\
&+ \frac{25N^6+213N^5+491N^4-1007N^3-7942N^2-15988N-10340}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_1(N)^2 \\
&+ \frac{-85N^6-1469N^5-8965N^4-23889N^3-25644N^2-3724N+5780}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_2(N) \\
&- \frac{2(94N^{10}+2202N^9+22629N^8+133916N^7+505769N^6+\dots+1817100N+563760)}{3(N+1)^2(N+2)^3(N+3)^3(N+4)^3(N+5)} S_1(N) \\
&- \left. \frac{2(44N^{11}+1696N^{10}+26555N^9+230482N^8+\dots+4371092N+623040)}{3(N+1)^3(N+2)^3(N+3)^3(N+4)^3(N+5)} \right) \varepsilon^{-1} \\
&+ \left(\dots \right) \varepsilon^0 \quad \text{Arising objects:} \\
&\quad \zeta_2, \zeta_3, (-1)^N, 2^N, S_{-3}(N), S_1(N), S_2(N), S_3(N), S_4(N), S_{-2,1}(N), \\
&\quad S_{2,1}(N), S_{3,1}(N)
\end{aligned}$$

[J.A.M. Vermaseren, 1998; J. Blümlein/S. Kurth, 1998]

$$\begin{aligned}
I_4(N) &= \left(\frac{64(N^2+N-1)}{3(N+1)(N+2)(N+3)(N+4)} - \frac{64S_1(N)}{3(N+3)(N+4)} \right) \varepsilon^{-3} \\
&+ \left(\frac{4(5N+27)}{3(N+2)(N+3)(N+4)} S_1(N)^2 - \frac{4(3N^5+68N^4+379N^3+648N^2-98N-696)}{3(N+1)(N+2)^2(N+3)^2(N+4)^2} S_1(N) \right. \\
&+ \left. \frac{4(14N^6+214N^5+1179N^4+3050N^3+4097N^2+3094N+1200)}{3(N+1)^2(N+2)^2(N+3)^2(N+4)^2} + \frac{4(N+1)(4N+17)S_2(N)}{3(N+2)(N+3)(N+4)} \right) \varepsilon^{-2} \\
&+ \left(\frac{-23N^2-35N-176}{9(N+2)(N+3)(N+4)(N+5)} S_1(N)^3 - \frac{2(10N^2+53N+106)}{3(N+2)(N+3)(N+4)} S_{2,1}(N) + \zeta_2 \left(\frac{8(N^2+N-1)}{(N+1)(N+2)(N+3)(N+4)} \right) \right. \\
&- \frac{8}{(N+3)(N+4)} S_1(N) + \frac{-8N^3-95N^2-171N-56}{3(N+2)(N+3)(N+4)(N+5)} S_2(N) S_1(N) + \frac{2(30N^3+469N^2+1873N+2542)}{9(N+2)(N+3)(N+4)(N+5)} S_3(N) \\
&+ \frac{25N^6+213N^5+491N^4-1007N^3-7942N^2-15988N-10340}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_1(N)^2 \\
&+ \frac{-85N^6-1469N^5-8965N^4-23889N^3-25644N^2-3724N+5780}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_2(N) \\
&- \frac{2(94N^{10}+2202N^9+22629N^8+133916N^7+505769N^6+\dots+1817100N+563760)}{3(N+1)^2(N+2)^3(N+3)^3(N+4)^3(N+5)} S_1(N) \\
&- \left. \frac{2(44N^{11}+1696N^{10}+26555N^9+230482N^8+\dots+4371092N+623040)}{3(N+1)^3(N+2)^3(N+3)^3(N+4)^3(N+5)} \right) \varepsilon^{-1} \\
&+ \left(\dots \right) \varepsilon^0 \quad \text{Arising objects:}
\end{aligned}$$

$$\begin{aligned}
&\zeta_2, \zeta_3, (-1)^N, 2^N, S_{-3}(N), S_1(N), S_2(N), S_3(N), S_4(N), S_{-2,1}(N), \\
&S_{2,1}(N), S_{3,1}(N), S_1\left(\frac{1}{2}, N\right), S_1(2, N), S_3\left(\frac{1}{2}, N\right), S_{1,1}\left(1, \frac{1}{2}, N\right), \\
&S_{1,1}\left(2, \frac{1}{2}, N\right), S_{2,1,1}(N), S_{2,1}\left(\frac{1}{2}, 1, N\right), S_{2,1}\left(1, \frac{1}{2}, N\right), S_{3,1}\left(\frac{1}{2}, 2, N\right), \\
&S_{1,1,1}\left(1, 1, \frac{1}{2}, N\right), S_{2,1,1}\left(1, \frac{1}{2}, 2, N\right), S_{1,1,1,1}\left(2, \frac{1}{2}, 1, 1, N\right)
\end{aligned}$$

$$\begin{aligned}
I_4(N) = & \left(\frac{64(N^2+N-1)}{3(N+1)(N+2)(N+3)(N+4)} - \frac{64S_1(N)}{3(N+3)(N+4)} \right) \varepsilon^{-3} \\
& + \left(\frac{4(5N+27)}{3(N+2)(N+3)(N+4)} S_1(N)^2 - \frac{4(3N^5+68N^4+379N^3+648N^2-98N-696)}{3(N+1)(N+2)^2(N+3)^2(N+4)^2} S_1(N) \right. \\
& + \left. \frac{4(14N^6+214N^5+1179N^4+3050N^3+4097N^2+3094N+1200)}{3(N+1)^2(N+2)^2(N+3)^2(N+4)^2} + \frac{4(N+1)(4N+17)S_2(N)}{3(N+2)(N+3)(N+4)} \right) \varepsilon^{-2} \\
& + \left(\frac{-23N^2-35N-176}{9(N+2)(N+3)(N+4)(N+5)} S_1(N)^3 - \frac{2(10N^2+53N+106)}{3(N+2)(N+3)(N+4)} S_{2,1}(N) + \zeta_2 \left(\frac{8(N^2+N-1)}{(N+1)(N+2)(N+3)(N+4)} \right. \right. \\
& - \left. \frac{8}{(N+3)(N+4)} S_1(N) \right) + \frac{-8N^3-95N^2-171N-56}{3(N+2)(N+3)(N+4)(N+5)} S_2(N)S_1(N) + \frac{2(30N^3+469N^2+1873N+2542)}{9(N+2)(N+3)(N+4)(N+5)} S_3(N) \\
& + \frac{25N^6+213N^5+491N^4-1007N^3-7942N^2-15988N-10340}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_1(N)^2 \\
& + \frac{-85N^6-1469N^5-8965N^4-23889N^3-25644N^2-3724N+5780}{3(N+1)(N+2)^2(N+3)^2(N+4)^2(N+5)} S_2(N) \\
& - \frac{2(94N^{10}+2202N^9+22629N^8+133916N^7+505769N^6+\dots+1817100N+563760)}{3(N+1)^2(N+2)^3(N+3)^3(N+4)^3(N+5)} S_1(N) \\
& - \left. \frac{2(44N^{11}+1696N^{10}+26555N^9+230482N^8+\dots+4371092N+623040)}{3(N+1)^3(N+2)^3(N+3)^3(N+4)^3(N+5)} \right) \varepsilon^{-1} \\
& + \left(\dots \right) \varepsilon^0 \quad \text{Arising objects:}
\end{aligned}$$

$$S_{1,1,1,1} \left(2, \frac{1}{2}, 1, 1, N \right) = \sum_{k=1}^N \frac{2^k \sum_{i=1}^k \frac{2^{-i} \sum_{j=1}^i \sum_{r=1}^j \frac{1}{r}}{j}}{k}$$

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.
- ▶ We obtained the ϵ -expansions of rather complicated master integrals.

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.
- ▶ We obtained the ϵ -expansions of rather complicated master integrals.
- ▶ Using these expansions we calculated easily the most complicated ladder graphs with 6 massive fermion lines
(using `Sigma.m`, `HarmonicSums.m`, `EvaluateMultiSums.m`, `SumProduction.m`).

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.
- ▶ We obtained the ϵ -expansions of rather complicated master integrals.
- ▶ Using these expansions we calculated easily the most complicated ladder graphs with 6 massive fermion lines
(using `Sigma.m`, `HarmonicSums.m`, `EvaluateMultiSums.m`, `SumProduction.m`).
- ▶ All ladder-topologies for 3-loop massive operator matrix elements can be calculated in this way.

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.
- ▶ We obtained the ϵ -expansions of rather complicated master integrals.
- ▶ Using these expansions we calculated easily the most complicated ladder graphs with 6 massive fermion lines
(using `Sigma.m`, `HarmonicSums.m`, `EvaluateMultiSums.m`, `SumProduction.m`).
- ▶ All ladder-topologies for 3-loop massive operator matrix elements can be calculated in this way.
- ▶ The mass production is ready for graphs depending on the same master integrals.

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.
- ▶ We obtained the ϵ -expansions of rather complicated master integrals.
- ▶ Using these expansions we calculated easily the most complicated ladder graphs with 6 massive fermion lines
(using `Sigma.m`, `HarmonicSums.m`, `EvaluateMultiSums.m`, `SumProduction.m`).
- ▶ All ladder-topologies for 3-loop massive operator matrix elements can be calculated in this way.
- ▶ The mass production is ready for graphs depending on the same master integrals.
- ▶ We used this technology for a few integrals emerging in the pure-singlet case.

Conclusion

- ▶ We presented a new method to solve coupled systems of differential and difference equations which emerge in massive Feynman diagram calculations.
- ▶ We obtained the ϵ -expansions of rather complicated master integrals.
- ▶ Using these expansions we calculated easily the most complicated ladder graphs with 6 massive fermion lines
(using `Sigma.m`, `HarmonicSums.m`, `EvaluateMultiSums.m`, `SumProduction.m`).
- ▶ All ladder-topologies for 3-loop massive operator matrix elements can be calculated in this way.
- ▶ The mass production is ready for graphs depending on the same master integrals.
- ▶ We used this technology for a few integrals emerging in the pure-singlet case.
- ▶ More involved massive 3-loop topologies are currently investigated.