

# Polynomial Reduction

Christoph Schwarzweiler  
University of Tuebingen

**Summary.** We continue the formalization of [8] towards Gröbner Bases. In this article we introduce reduction of polynomials and prove its termination, its adequateness for ideal congruence as well as the translation lemma used later to show confluence of reduction.

MML Identifier: POLYRED.

WWW: <http://mizar.org/JFM/Vol14/polyred.html>

The articles [22], [21], [27], [12], [28], [30], [29], [10], [11], [4], [3], [17], [6], [23], [13], [5], [25], [2], [7], [26], [9], [16], [14], [19], [1], [24], [18], [15], and [20] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

Let  $n$  be an ordinal number and let  $R$  be a non trivial zero structure. One can verify that there exists a monomial of  $n, R$  which is non-zero.

One can verify that there exists a field which is non trivial.

Let us observe that every left zeroed add-right-cancelable right distributive left unital commutative associative non empty double loop structure which is field-like is also integral domain-like.

Let  $n$  be an ordinal number, let  $L$  be an add-associative right complementable left zeroed right zeroed unital distributive integral domain-like non trivial double loop structure, and let  $p, q$  be non-zero finite-Support series of  $n, L$ . One can check that  $p * q$  is non-zero.

## 2. MORE ON POLYNOMIALS AND MONOMIALS

The following propositions are true:

- (1) Let  $X$  be a set,  $L$  be an Abelian add-associative right zeroed right complementable non empty loop structure, and  $p, q$  be series of  $X, L$ . Then  $-(p + q) = -p + -q$ .
- (2) For every set  $X$  and for every left zeroed non empty loop structure  $L$  and for every series  $p$  of  $X, L$  holds  $0_X L + p = p$ .
- (3) Let  $X$  be a set,  $L$  be an add-associative right zeroed right complementable non empty loop structure, and  $p$  be a series of  $X, L$ . Then  $-p + p = 0_X L$  and  $p + -p = 0_X L$ .
- (4) Let  $n$  be a set,  $L$  be an add-associative right zeroed right complementable non empty loop structure, and  $p$  be a series of  $n, L$ . Then  $p - 0_n L = p$ .
- (5) Let  $n$  be an ordinal number,  $L$  be an add-associative right complementable right zeroed add-left-cancelable left distributive non empty double loop structure, and  $p$  be a series of  $n, L$ . Then  $0_n L * p = 0_n L$ .

- (6) Let  $n$  be an ordinal number,  $L$  be an Abelian right zeroed add-associative right complementable unital distributive associative commutative non trivial double loop structure, and  $p, q$  be polynomials of  $n, L$ . Then  $-p * q = (-p) * q$  and  $-p * q = p * -q$ .
- (7) Let  $n$  be an ordinal number,  $L$  be an add-associative right complementable right zeroed distributive non empty double loop structure,  $p$  be a polynomial of  $n, L$ ,  $m$  be a monomial of  $n, L$ , and  $b$  be a bag of  $n$ . Then  $(m * p)(\text{term } m + b) = m(\text{term } m) \cdot p(b)$ .
- (8) Let  $X$  be a set,  $L$  be a right zeroed add-left-cancelable left distributive non empty double loop structure, and  $p$  be a series of  $X, L$ . Then  $0_L \cdot p = 0_X L$ .
- (9) Let  $X$  be a set,  $L$  be an add-associative right zeroed right complementable distributive non empty double loop structure,  $p$  be a series of  $X, L$ , and  $a$  be an element of  $L$ . Then  $-a \cdot p = (-a) \cdot p$  and  $-a \cdot p = a \cdot -p$ .
- (10) Let  $X$  be a set,  $L$  be a left distributive non empty double loop structure,  $p$  be a series of  $X, L$ , and  $a, a'$  be elements of  $L$ . Then  $a \cdot p + a' \cdot p = (a + a') \cdot p$ .
- (11) Let  $X$  be a set,  $L$  be an associative non empty multiplicative loop with zero structure,  $p$  be a series of  $X, L$ , and  $a, a'$  be elements of  $L$ . Then  $(a \cdot a') \cdot p = a \cdot (a' \cdot p)$ .
- (12) Let  $n$  be an ordinal number,  $L$  be an add-associative right zeroed right complementable unital associative commutative distributive non empty double loop structure,  $p, p'$  be series of  $n, L$ , and  $a$  be an element of  $L$ . Then  $a \cdot (p * p') = p * (a \cdot p')$ .

### 3. MULTIPLICATION OF POLYNOMIALS WITH BAGS

Let  $n$  be an ordinal number, let  $b$  be a bag of  $n$ , let  $L$  be a non empty zero structure, and let  $p$  be a series of  $n, L$ . The functor  $b * p$  yielding a series of  $n, L$  is defined by:

- (Def. 1) For every bag  $b'$  of  $n$  such that  $b \mid b'$  holds  $(b * p)(b') = p(b' -' b)$  and for every bag  $b'$  of  $n$  such that  $b \nmid b'$  holds  $(b * p)(b') = 0_L$ .

Let  $n$  be an ordinal number, let  $b$  be a bag of  $n$ , let  $L$  be a non empty zero structure, and let  $p$  be a finite-Support series of  $n, L$ . Note that  $b * p$  is finite-Support.

Next we state a number of propositions:

- (13) Let  $n$  be an ordinal number,  $b, b'$  be bags of  $n, L$  be a non empty zero structure, and  $p$  be a series of  $n, L$ . Then  $(b * p)(b' + b) = p(b')$ .
- (14) Let  $n$  be an ordinal number,  $L$  be a non empty zero structure,  $p$  be a polynomial of  $n, L$ , and  $b$  be a bag of  $n$ . Then  $\text{Support}(b * p) \subseteq \{b + b'; b' \text{ ranges over elements of } \text{Bags } n : b' \in \text{Support } p\}$ .
- (15) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n, L$  be a non trivial zero structure,  $p$  be a non-zero polynomial of  $n, L$ , and  $b$  be a bag of  $n$ . Then  $\text{HT}(b * p, T) = b + \text{HT}(p, T)$ .
- (16) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n, L$  be a non empty zero structure,  $p$  be a polynomial of  $n, L$ , and  $b, b'$  be bags of  $n$ . If  $b' \in \text{Support}(b * p)$ , then  $b' \leq_T b + \text{HT}(p, T)$ .
- (17) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty zero structure, and  $p$  be a series of  $n, L$ . Then  $\text{EmptyBag } n * p = p$ .
- (18) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty zero structure,  $p$  be a series of  $n, L$ , and  $b_1, b_2$  be bags of  $n$ . Then  $(b_1 + b_2) * p = b_1 * (b_2 * p)$ .
- (19) Let  $n$  be an ordinal number,  $L$  be an add-associative right zeroed right complementable distributive non trivial double loop structure,  $p$  be a polynomial of  $n, L$ , and  $a$  be an element of  $L$ . Then  $\text{Support}(a \cdot p) \subseteq \text{Support } p$ .

- (20) Let  $n$  be an ordinal number,  $L$  be an integral domain-like non trivial double loop structure,  $p$  be a polynomial of  $n, L$ , and  $a$  be a non-zero element of  $L$ . Then  $\text{Support } p \subseteq \text{Support}(a \cdot p)$ .
- (21) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be an add-associative right zeroed right complementable distributive integral domain-like non trivial double loop structure,  $p$  be a polynomial of  $n, L$ , and  $a$  be a non-zero element of  $L$ . Then  $\text{HT}(a \cdot p, T) = \text{HT}(p, T)$ .
- (22) Let  $n$  be an ordinal number,  $L$  be an add-associative right complementable right zeroed distributive non trivial double loop structure,  $p$  be a series of  $n, L$ ,  $b$  be a bag of  $n$ , and  $a$  be an element of  $L$ . Then  $a \cdot (b * p) = \text{Monom}(a, b) * p$ .
- (23) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n, L$  be an add-associative right complementable right zeroed unital distributive integral domain-like non trivial double loop structure,  $p$  be a non-zero polynomial of  $n, L$ ,  $q$  be a polynomial of  $n, L$ , and  $m$  be a non-zero monomial of  $n, L$ . If  $\text{HT}(p, T) \in \text{Support } q$ , then  $\text{HT}(m * p, T) \in \text{Support}(m * q)$ .

#### 4. ORDERS ON POLYNOMIALS

Let  $n$  be an ordinal number and let  $T$  be a connected term order of  $n$ . Observe that  $\langle \text{Bags } n, T \rangle$  is connected.

Let  $n$  be a natural number and let  $T$  be an admissible term order of  $n$ . Note that  $\langle \text{Bags } n, T \rangle$  is well founded.

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be a non empty zero structure, and let  $p, q$  be polynomials of  $n, L$ . The predicate  $p \leq_T q$  is defined by:

(Def. 2)  $\langle \text{Support } p, \text{Support } q \rangle \in \text{FinOrd}(\langle \text{Bags } n, T \rangle)$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be a non empty zero structure, and let  $p, q$  be polynomials of  $n, L$ . The predicate  $p <_T q$  is defined as follows:

(Def. 3)  $p \leq_T q$  and  $\text{Support } p \neq \text{Support } q$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be a non empty zero structure, and let  $p$  be a polynomial of  $n, L$ . The functor  $\text{Support}(p, T)$  yielding an element of  $\text{Fin}$  (the carrier of  $\langle \text{Bags } n, T \rangle$ ) is defined as follows:

(Def. 4)  $\text{Support}(p, T) = \text{Support } p$ .

We now state a number of propositions:

- (24) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non trivial zero structure, and  $p$  be a non-zero polynomial of  $n, L$ . Then  $\text{PosetMax } \text{Support}(p, T) = \text{HT}(p, T)$ .
- (25) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty loop structure, and  $p$  be a polynomial of  $n, L$ . Then  $p \leq_T p$ .
- (26) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty loop structure, and  $p, q$  be polynomials of  $n, L$ . Then  $p \leq_T q$  and  $q \leq_T p$  if and only if  $\text{Support } p = \text{Support } q$ .
- (27) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty loop structure, and  $p, q, r$  be polynomials of  $n, L$ . If  $p \leq_T q$  and  $q \leq_T r$ , then  $p \leq_T r$ .
- (28) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty loop structure, and  $p, q$  be polynomials of  $n, L$ . Then  $p \leq_T q$  or  $q \leq_T p$ .
- (29) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be a non empty loop structure, and  $p, q$  be polynomials of  $n, L$ . Then  $p \leq_T q$  if and only if  $q \not<_T p$ .

- (30) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be a non empty zero structure, and  $p$  be a polynomial of  $n, L$ . Then  $0_n L \leq_T p$ .
- (31) Let  $n$  be a natural number,  $T$  be an admissible connected term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed unital distributive non trivial double loop structure, and  $P$  be a non empty subset of  $\text{Polynom-Ring}(n, L)$ . Then there exists a polynomial  $p$  of  $n, L$  such that  $p \in P$  and for every polynomial  $q$  of  $n, L$  such that  $q \in P$  holds  $p \leq_T q$ .
- (32) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed non trivial loop structure, and  $p, q$  be polynomials of  $n, L$ . Then  $p <_T q$  if and only if one of the following conditions is satisfied:
- (i)  $p = 0_n L$  and  $q \neq 0_n L$ , or
  - (ii)  $\text{HT}(p, T) <_T \text{HT}(q, T)$ , or
  - (iii)  $\text{HT}(p, T) = \text{HT}(q, T)$  and  $\text{Red}(p, T) <_T \text{Red}(q, T)$ .
- (33) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed non trivial loop structure, and  $p$  be a non-zero polynomial of  $n, L$ . Then  $\text{Red}(p, T) <_T \text{HM}(p, T)$ .
- (34) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed non trivial loop structure, and  $p$  be a polynomial of  $n, L$ . Then  $\text{HM}(p, T) \leq_T p$ .
- (35) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed non trivial loop structure, and  $p$  be a non-zero polynomial of  $n, L$ . Then  $\text{Red}(p, T) <_T p$ .

## 5. POLYNOMIAL REDUCTION

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, let  $f, p, g$  be polynomials of  $n, L$ , and let  $b$  be a bag of  $n$ . We say that  $f$  reduces to  $g, p, b, T$  if and only if:

- (Def. 5)  $f \neq 0_n L$  and  $p \neq 0_n L$  and  $b \in \text{Support } f$  and there exists a bag  $s$  of  $n$  such that  $s + \text{HT}(p, T) = b$  and  $g = f - \frac{f(b)}{\text{HC}(p, T)} \cdot (s * p)$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, and let  $f, p, g$  be polynomials of  $n, L$ . We say that  $f$  reduces to  $g, p, T$  if and only if:

- (Def. 6) There exists a bag  $b$  of  $n$  such that  $f$  reduces to  $g, p, b, T$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, let  $f, g$  be polynomials of  $n, L$ , and let  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ . We say that  $f$  reduces to  $g, P, T$  if and only if:

- (Def. 7) There exists a polynomial  $p$  of  $n, L$  such that  $p \in P$  and  $f$  reduces to  $g, p, T$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, and let  $f, p$  be polynomials of  $n, L$ . We say that  $f$  is reducible wrt  $p, T$  if and only if:

- (Def. 8) There exists a polynomial  $g$  of  $n, L$  such that  $f$  reduces to  $g, p, T$ .

We introduce  $f$  is irreducible wrt  $p, T$  and  $f$  is in normal form wrt  $p, T$  as antonyms of  $f$  is reducible wrt  $p, T$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, let  $f$  be a polynomial of  $n, L$ , and let  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ . We say that  $f$  is reducible wrt  $P, T$  if and only if:

(Def. 9) There exists a polynomial  $g$  of  $n, L$  such that  $f$  reduces to  $g, P, T$ .

We introduce  $f$  is irreducible wrt  $P, T$  and  $f$  is in normal form wrt  $P, T$  as antonyms of  $f$  is reducible wrt  $P, T$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, and let  $f, p, g$  be polynomials of  $n, L$ . We say that  $f$  top reduces to  $g, p, T$  if and only if:

(Def. 10)  $f$  reduces to  $g, p, \text{HT}(f, T), T$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, and let  $f, p$  be polynomials of  $n, L$ . We say that  $f$  is top reducible wrt  $p, T$  if and only if:

(Def. 11) There exists a polynomial  $g$  of  $n, L$  such that  $f$  top reduces to  $g, p, T$ .

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, let  $f$  be a polynomial of  $n, L$ , and let  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ . We say that  $f$  is top reducible wrt  $P, T$  if and only if:

(Def. 12) There exists a polynomial  $p$  of  $n, L$  such that  $p \in P$  and  $f$  is top reducible wrt  $p, T$ .

Next we state several propositions:

- (36) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure,  $f$  be a polynomial of  $n, L$ , and  $p$  be a non-zero polynomial of  $n, L$ . Then  $f$  is reducible wrt  $p, T$  if and only if there exists a bag  $b$  of  $n$  such that  $b \in \text{Support } f$  and  $\text{HT}(p, T) \mid b$ .
- (37) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, and  $p$  be a polynomial of  $n, L$ . Then  $0_n L$  is irreducible wrt  $p, T$ .
- (38) Let  $n$  be an ordinal number,  $T$  be an admissible connected term order of  $n, L$  be an add-associative right complementable right zeroed commutative associative well unital distributive Abelian field-like non degenerated non empty double loop structure,  $f, p$  be polynomials of  $n, L$ , and  $m$  be a non-zero monomial of  $n, L$ . If  $f$  reduces to  $f - m * p, p, T$ , then  $\text{HT}(m * p, T) \in \text{Support } f$ .
- (39) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n, L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure,  $f, p, g$  be polynomials of  $n, L$ , and  $b$  be a bag of  $n$ . If  $f$  reduces to  $g, p, b, T$ , then  $b \notin \text{Support } g$ .
- (40) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n, L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure,  $f, p, g$  be polynomials of  $n, L$ , and  $b, b'$  be bags of  $n$ . Suppose  $b <_T b'$ . If  $f$  reduces to  $g, p, b, T$ , then  $b' \in \text{Support } g$  iff  $b' \in \text{Support } f$ .

- (41) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure,  $f, p, g$  be polynomials of  $n, L$ , and  $b, b'$  be bags of  $n$ . If  $b <_T b'$ , then if  $f$  reduces to  $g, p, b, T$ , then  $f(b') = g(b')$ .
- (42) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure, and  $f, p, g$  be polynomials of  $n, L$ . Suppose  $f$  reduces to  $g, p, T$ . Let  $b$  be a bag of  $n$ . If  $b \in \text{Support } g$ , then  $b \leq_T \text{HT}(f, T)$ .
- (43) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure, and  $f, p, g$  be polynomials of  $n, L$ . If  $f$  reduces to  $g, p, T$ , then  $g <_T f$ .

## 6. POLYNOMIAL REDUCTION RELATION

Let  $n$  be an ordinal number, let  $T$  be a connected term order of  $n$ , let  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure, and let  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ . The functor  $\text{PolyRedRel}(P, T)$  yielding a relation between (the carrier of  $\text{Polynom-Ring}(n, L) \setminus \{0_n L\}$  and the carrier of  $\text{Polynom-Ring}(n, L)$ ) is defined as follows:

(Def. 13) For all polynomials  $p, q$  of  $n, L$  holds  $\langle p, q \rangle \in \text{PolyRedRel}(P, T)$  iff  $p$  reduces to  $q, P, T$ .

One can prove the following proposition

- (44) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure,  $f, g$  be polynomials of  $n, L$ , and  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ . If  $\text{PolyRedRel}(P, T)$  reduces  $f$  to  $g$ , then  $g \leq_T f$  but  $g = 0_n L$  or  $\text{HT}(g, T) \leq_T \text{HT}(f, T)$ .

Let  $n$  be a natural number, let  $T$  be a connected admissible term order of  $n$ , let  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure, and let  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ . Observe that  $\text{PolyRedRel}(P, T)$  is strongly-normalizing.

Next we state several propositions:

- (45) Let  $n$  be a natural number,  $T$  be an admissible connected term order of  $n$ ,  $L$  be an add-associative right complementable left zeroed right zeroed commutative associative well unital distributive Abelian field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, h$  be polynomials of  $n, L$ . If  $f \in P$ , then  $\text{PolyRedRel}(P, T)$  reduces  $h * f$  to  $0_n L$ .
- (46) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ ,  $f, g$  be polynomials of  $n, L$ , and  $m$  be a non-zero monomial of  $n, L$ . If  $f$  reduces to  $g, P, T$ , then  $m * f$  reduces to  $m * g, P, T$ .
- (47) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ ,  $f, g$  be polynomials of  $n, L$ , and  $m$  be a monomial of  $n, L$ . If  $\text{PolyRedRel}(P, T)$  reduces  $f$  to  $g$ , then  $\text{PolyRedRel}(P, T)$  reduces  $m * f$  to  $m * g$ .

- (48) Let  $n$  be an ordinal number,  $T$  be a connected admissible term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ ,  $f$  be a polynomial of  $n, L$ , and  $m$  be a monomial of  $n, L$ . If  $\text{PolyRedRel}(P, T)$  reduces  $f$  to  $0_n L$ , then  $\text{PolyRedRel}(P, T)$  reduces  $m * f$  to  $0_n L$ .
- (49) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive Abelian field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, g, h, h_1$  be polynomials of  $n, L$ . Suppose  $f - g = h$  and  $\text{PolyRedRel}(P, T)$  reduces  $h$  to  $h_1$ . Then there exist polynomials  $f_1, g_1$  of  $n, L$  such that  $f_1 - g_1 = h_1$  and  $\text{PolyRedRel}(P, T)$  reduces  $f$  to  $f_1$  and  $\text{PolyRedRel}(P, T)$  reduces  $g$  to  $g_1$ .
- (50) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive Abelian field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, g$  be polynomials of  $n, L$ . Suppose  $\text{PolyRedRel}(P, T)$  reduces  $f - g$  to  $0_n L$ . Then  $f$  and  $g$  are convergent w.r.t.  $\text{PolyRedRel}(P, T)$ .
- (51) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an add-associative right complementable right zeroed commutative associative well unital distributive Abelian field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, g$  be polynomials of  $n, L$ . Suppose  $\text{PolyRedRel}(P, T)$  reduces  $f - g$  to  $0_n L$ . Then  $f$  and  $g$  are convertible w.r.t.  $\text{PolyRedRel}(P, T)$ .

Let  $R$  be a non empty loop structure, let  $I$  be a subset of  $R$ , and let  $a, b$  be elements of  $R$ . The predicate  $a \equiv b(\text{mod } I)$  is defined by:

(Def. 14)  $a - b \in I$ .

Next we state several propositions:

- (52) Let  $R$  be an add-associative left zeroed right zeroed right complementable right distributive non empty double loop structure,  $I$  be a right ideal non empty subset of  $R$ , and  $a$  be an element of  $R$ . Then  $a \equiv a(\text{mod } I)$ .
- (53) Let  $R$  be an add-associative right zeroed right complementable right unital right distributive non empty double loop structure,  $I$  be a right ideal non empty subset of  $R$ , and  $a, b$  be elements of  $R$ . If  $a \equiv b(\text{mod } I)$ , then  $b \equiv a(\text{mod } I)$ .
- (54) Let  $R$  be an add-associative right zeroed right complementable non empty loop structure,  $I$  be an add closed non empty subset of  $R$ , and  $a, b, c$  be elements of  $R$ . If  $a \equiv b(\text{mod } I)$  and  $b \equiv c(\text{mod } I)$ , then  $a \equiv c(\text{mod } I)$ .
- (55) Let  $R$  be an Abelian add-associative right zeroed right complementable unital distributive associative non trivial double loop structure,  $I$  be an add closed non empty subset of  $R$ , and  $a, b, c, d$  be elements of  $R$ . If  $a \equiv b(\text{mod } I)$  and  $c \equiv d(\text{mod } I)$ , then  $a + c \equiv b + d(\text{mod } I)$ .
- (56) Let  $R$  be an add-associative right zeroed right complementable commutative distributive non empty double loop structure,  $I$  be an add closed right ideal non empty subset of  $R$ , and  $a, b, c, d$  be elements of  $R$ . If  $a \equiv b(\text{mod } I)$  and  $c \equiv d(\text{mod } I)$ , then  $a \cdot c \equiv b \cdot d(\text{mod } I)$ .
- (57) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, g$  be elements of  $\text{Polynom-Ring}(n, L)$ . If  $f$  and  $g$  are convertible w.r.t.  $\text{PolyRedRel}(P, T)$ , then  $f \equiv g(\text{mod } P\text{-ideal})$ .

- (58) Let  $n$  be a natural number,  $T$  be an admissible connected term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non degenerated non empty double loop structure,  $P$  be a non empty subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, g$  be elements of  $\text{Polynom-Ring}(n, L)$ . If  $f \equiv g \pmod{P\text{-ideal}}$ , then  $f$  and  $g$  are convertible w.r.t.  $\text{PolyRedRel}(P, T)$ .
- (59) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f, g$  be polynomials of  $n, L$ . If  $\text{PolyRedRel}(P, T)$  reduces  $f$  to  $g$ , then  $f - g \in P\text{-ideal}$ .
- (60) Let  $n$  be an ordinal number,  $T$  be a connected term order of  $n$ ,  $L$  be an Abelian add-associative right complementable right zeroed commutative associative well unital distributive field-like non trivial double loop structure,  $P$  be a subset of  $\text{Polynom-Ring}(n, L)$ , and  $f$  be a polynomial of  $n, L$ . If  $\text{PolyRedRel}(P, T)$  reduces  $f$  to  $0_n L$ , then  $f \in P\text{-ideal}$ .

## REFERENCES

- [1] Jonathan Backer, Piotr Rudnicki, and Christoph Schwarzeweller. Ring ideals. *Journal of Formalized Mathematics*, 12, 2000. [http://mizar.org/JFM/Vol12/ideal\\_1.html](http://mizar.org/JFM/Vol12/ideal_1.html).
- [2] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/card\\_1.html](http://mizar.org/JFM/Vol1/card_1.html).
- [3] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/nat\\_1.html](http://mizar.org/JFM/Vol1/nat_1.html).
- [4] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [5] Grzegorz Bancerek. Reduction relations. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/rewritel.html>.
- [6] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/waybel\\_0.html](http://mizar.org/JFM/Vol8/waybel_0.html).
- [7] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finseq\\_1.html](http://mizar.org/JFM/Vol1/finseq_1.html).
- [8] Thomas Becker and Volker Weispfenning. *Gröbner Bases: A Computational Approach to Commutative Algebra*. Springer-Verlag, New York, Berlin, 1993.
- [9] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/realset1.html>.
- [10] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [11] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_2.html](http://mizar.org/JFM/Vol1/funct_2.html).
- [12] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/zfmisc\\_1.html](http://mizar.org/JFM/Vol1/zfmisc_1.html).
- [13] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finset\\_1.html](http://mizar.org/JFM/Vol1/finset_1.html).
- [14] Eugeniusz Kusak, Wojciech Leńczuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/vectsp\\_1.html](http://mizar.org/JFM/Vol1/vectsp_1.html).
- [15] Gilbert Lee and Piotr Rudnicki. On ordering of bags. *Journal of Formalized Mathematics*, 14, 2002. <http://mizar.org/JFM/Vol14/bagorder.html>.
- [16] Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/vectsp\\_2.html](http://mizar.org/JFM/Vol2/vectsp_2.html).
- [17] Michał Muzalewski and Wojciech Skaba. From loops to abelian multiplicative groups with zero. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/algstr\\_1.html](http://mizar.org/JFM/Vol2/algstr_1.html).
- [18] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/polynom1.html>.
- [19] Christoph Schwarzeweller. More on multivariate polynomials: Monomials and constant polynomials. *Journal of Formalized Mathematics*, 13, 2001. <http://mizar.org/JFM/Vol13/polynom7.html>.
- [20] Christoph Schwarzeweller. Term orders. *Journal of Formalized Mathematics*, 14, 2002. <http://mizar.org/JFM/Vol14/termord.html>.



- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [22] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [23] Andrzej Trybulec and Agata Darmochwał. Boolean domains. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finsub\\_1.html](http://mizar.org/JFM/Vol1/finsub_1.html).
- [24] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/orders\\_1.html](http://mizar.org/JFM/Vol1/orders_1.html).
- [25] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/rlvect\\_1.html](http://mizar.org/JFM/Vol1/rlvect_1.html).
- [26] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/finseq\\_4.html](http://mizar.org/JFM/Vol2/finseq_4.html).
- [27] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [28] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).
- [29] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relset\\_1.html](http://mizar.org/JFM/Vol1/relset_1.html).
- [30] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_2.html](http://mizar.org/JFM/Vol1/relat_2.html).

*Received December 20, 2002*

*Published January 2, 2004*

---